arXiv:1106.0742v2 [math.AC] 22 Sep 2011

# REES ALGEBRAS OF DIAGONAL IDEALS

KUEI-NUAN LIN

Abstract. There is a natural epimorphism from the symmetric algebra to the Rees algebra of an ideal. When this epimorphism is an isomorphism, we say that the ideal is of linear type. Given two determinantal rings over a field, we consider the diagonal ideal, the kernel of the multiplication map. We prove that the diagonal ideal is of linear type and recover the defining ideal of the Rees algebra in some special cases. The special fiber ring of the diagonal ideal is the homogeneous coordinate ring of the join variety.

## 1. INTRODUCTION

In this paper we address the problem of determining the equations that define the Rees algebra of an ideal. Besides encoding asymptotic properties of the powers of an ideal, the Rees algebra realizes, algebraically, the blow-up of a variety along a subvariety. Though blowing up is a fundamental operation in the birational study of algebraic varieties and, in particular, in the process of desingularization, an explicit description of the resulting variety in terms of defining equations remains a difficult problem.

Let $I$ be an ideal in a Noetherian ring $R$. The Rees algebra $\mathcal{R}(I)$ of $I$ is the graded subalgebra $R[It] \cong \oplus_{n\geq 0} I^n$ of $R[t]$. When $I$ is generated by $f_1, ..., f_u$, there is a natural map $\phi$ from $R[t_1, ..., t_u]$ to $\mathcal{R}(I)$ sending $t_i$ to $f_i t$. The kernel of $\phi$ is the defining ideal of $\mathcal{R}(I)$ in the ring $R[t_1, ..., t_u]$. There is another natural map $\psi$ from $\mathrm{Sym}(R^u) = R[t_1, ..., t_u]$ to $\mathrm{Sym}(I)$, the symmetric algebra of $I$, and the kernel of $\psi$ is the defining ideal of $\mathrm{Sym}(I)$. This ideal is generated by the entries of the product of $(t_1, ..., t_u)$ and the presentation matrix of $I$. The defining ideal of $\mathrm{Sym}(I)$ is contained in the kernel of $\phi$. Hence we have a surjective map from $\mathrm{Sym}(I)$ to $\mathcal{R}(I)$. The ideal $I$ is said to be of *linear type* if $\mathrm{Sym}(I)$ is naturally isomorphic to $\mathcal{R}(I)$. Hence we obtain the defining equations of $\mathcal{R}(I)$ for free in this case.

In general, an ideal is not of linear type. The first known class of ideals of linear type are complete intersection ideals [M]. Ideals generated by $d$-sequences are another large class of ideals of linear type [H1], [V]. These sequences play a role in the theory of approximation complexes similar to the role regular sequences play

---

2000 *Mathematics Subject Classification.* Primary 13C40, 14M12 ; Secondary 13P10, 14Q15.

*Key words and phrases.* Rees Algebra, Join Variety, Determinantal Ring, Gröbner Basis, Symmetric Algebra.

in the theory of Koszul complexes. Later Herzog-Simis-Vansconcelos and Herzog-Vansconcelos-Villarreal used strongly Cohen-Macaulay and sliding depth conditions to describe classes of ideals of linear type [HSV1], [HSV2], [HVV]. Huneke proved that if $X$ is a generic $n \times n$ matrix and $I$ is the ideal of $n-1$ size minors of $X$ in $R = \mathbb{Z}[x_{ij}]$, then $I$ is of linear type [H2]. Villarreal showed the edge ideals of a tree or a graph with a unique odd cycle are ideals of linear type [Vi]. In this paper, we give a new class of ideals of linear type, diagonal ideals of determinantal rings.

Now we describe the setting of this work. Let $k$ be a field, $R$ be a polynomial ring over the field $k$ with variables $\{x_{ij}\}$, and $X$ is the generic $m \times n$ matrix $(x_{ij})$. Given two homogeneous $R$-ideals, $I_1$ and $I_2$, we consider the kernel of the multiplication map from $S = R/I_1 \otimes R/I_2$ to $R/(I_1 + I_2)$. The kernel is the *diagonal* ideal $\mathbb{D}$ of the ring $S$ and $\mathbb{D}$ is generated by the images of $x_i \otimes 1 - 1 \otimes x_i$ in the ring $S$. The main result of this paper shows that the ideal $\mathbb{D}$ is of linear type if $I_1$, $I_2$ are the ideals of maximal minors of submatrics of $X$. Notice $I_1$ and $I_2$ are in general not of linear type (see [H2], 2.6).

In this particular case, the *special fiber ring* of $I$, $\mathcal{F}(\mathbb{D}) = \mathcal{R}(\mathbb{D}) \otimes_S k$, is the homogeneous coordinate ring of the embedded join varieties of $V(I_1)$ and $V(I_2)$ in projective space $\mathbb{P}_k^{m \times n - 1}$. Hence when $\mathbb{D}$ is an ideal of linear type, the embedded join is the whole space. But it is not true in general that if the embedded join variety is the whole space, the diagonal ideal $\mathbb{D}$ is of linear type. See Example 2.2 in Section 2.

The proof of the main result is in Section 2. We now describe the idea of the proof. We use the defining ideals of $\mathrm{Sym}(\mathbb{D})$ to understand the defining ideals of $\mathcal{R}(\mathbb{D})$. We identify some specific equations in the defining ideal $\mathcal{J}$ of $\mathrm{Sym}(\mathbb{D})$, and consider the subideal $\mathcal{L}$ of $\mathcal{J}$ they generate.

Notice that $\mathcal{L} \subset \mathcal{J} \subset \mathcal{K}$, where $\mathcal{K}$ is the defining ideal of $\mathcal{R}(\mathbb{D})$, hence the goal is to prove that $\mathcal{L} = \mathcal{K}$. We use Buchberger's Algorithm to find a Gröbner basis of the ideal $\mathcal{L}$ with respect to some monomial order. We find a set of polynomials that are in the ideal $\mathcal{L}$ and show that all the remainders between elements in this set are zero. Hence we find a Gröbner basis of the ideal $\mathcal{L}$. Once we have the Gröbner basis, we have the generating set for the initial ideal $\mathrm{in}(\mathcal{L})$ of $\mathcal{L}$. This way we find a non zero-divisor modulo $\mathcal{L}$, which we may invert thereby reducing to the case of a smaller matrix. Thus we show that $\mathcal{L} = \mathcal{K}$. As a consequence, the two algebras $\mathrm{Sym}(\mathbb{D})$ and $\mathcal{R}(\mathbb{D})$ are naturally isomorphic and we obtain an explicit description of the defining equations of $\mathcal{R}(\mathbb{D})$.

**Acknowledgments:** This work is based on author's Ph. D. thesis from Purdue University under the direction of Professor Bernd Ulrich. The author is very grateful for so many useful suggestions from Professor Ulrich.

## 2. Main results

Let $k$ be a field, $2 \leq m \leq n$ integers, $X_{mn} = [x_{ij}]$, $Y_{mn} = [y_{ij}]$, $Z_{mn} = [z_{ij}]$, $m \times n$ matrices of variables over $k$. Let $2 \leq s_i \leq t_i$ integers, and let $X_{s_1 t_1}$, $Y_{s_2 t_2}$ be the submatrices of $X$ and $Y$ consisting of the first $s_i$ rows and first $t_i$ columns. We write $I = I_{s_1}(X_{s_1 t_1})$, $J = I_{s_2}(X_{s_2 t_2})$ the ideals of $k[X]$ generated by the maximal minors of $X_{s_1 t_1}$ and the maximal minors of $X_{s_2 t_2}$. Let $R_1 = k[X]/I$, $R_2 = k[X]/J$ be the two determinantal rings. We consider the diagonal ideal $\mathbb{D}$ of $R_1 \otimes_k R_2$, defined via the exact sequence

$$0 \longrightarrow \mathbb{D} \longrightarrow R_1 \otimes_k R_2 \xrightarrow{\text{mult.}} k[X]/(I+J) \longrightarrow 0.$$

The ideal $\mathbb{D}$ is generated by the images of $x_{ij} \otimes 1 - 1 \otimes x_{ij}$ in $R_1 \otimes_k R_2$.

We write the diagonal ideal $\mathbb{D} = (\{x_{ij} - y_{ij}\})$ in

$$S = k[X_{mn}, Y_{mn}]/(I_{s_1}(X_{s_1 t_1}), I_{s_2}(Y_{s_2 t_2})) \cong R_1 \otimes_k R_2.$$

We have a presentation of $\mathbb{D}$,

$$S^l \quad \xrightarrow{\phi} \quad S^{mn} \quad \longrightarrow \mathbb{D} \quad \longrightarrow \quad 0$$

From this we obtain a presentation of the symmetric algebra of $\mathbb{D}$,

$$0 \to (\text{image}(\phi)) = J \longrightarrow \text{Sym}(S^{mn}) = S[Z_{mn}] = T \longrightarrow \text{Sym}(\mathbb{D}) \to 0.$$

Here $J$ is the ideal generated by the entries of the row vector $[z_{11}, z_{12}, ..., z_{1n}, ...., z_{mn}] \cdot \phi$. Hence

$$\text{Sym}(\mathbb{D}) \cong T/J$$

where $J$ is generated by linear forms in the variables $z_{ij}$. We write $\mathcal{R}(\mathbb{D}) = T/K$, $J \subset K$. In general $K$ is not generated by linear forms. We can rewrite $\text{Sym}(\mathbb{D}) = T/J = k[X_{mn}, Y_{mn,} Z_{mn}]/\mathcal{J}$ and $\mathcal{R}(\mathbb{D}) = k[X_{mn}, Y_{mn}, Z_{mn}]/\mathcal{K}$. In this particular case, the *special fiber ring* of $I$, $\mathcal{F}(\mathbb{D}) = \mathcal{R}(\mathbb{D}) \otimes_S k$, is the homogeneous coordinate ring of the embedded join varieties of $V(I_1)$ and $V(I_2)$ in projective space $\mathbb{P}_k^{m \times n - 1}$.

**Theorem 2.1.** *The ideal $\mathbb{D}$ is of linear type if $I_1$ and $I_2$ are generated by the maximal minors of submatrices of $X$. So $\mathcal{R}(\mathbb{D}) \cong \text{Sym}(\mathbb{D})$.*

Hence the embedded join is the whole space in the above case. The following example showing that even the fiber ring is the whole space, the ideal $\mathbb{D}$ may not be of linear type in general.

**Example 2.2.** Let $X$, $Y$, and $Z$ be $3 \times 3$ matrices and $I_1 = I_3(X)$, $I_2 = I_2(X)$ be the ideal generated by $3 \times 3$ and $2 \times 2$ minors of $X$. Write

$$S = k[X, Y]/(I_3(X), I_2(Y)) \cong R_1 \otimes_k R_2,$$

$\mathrm{Sym}(\mathbb{D}) = S[Z]/J$ and $\mathcal{R}(\mathbb{D}) = S[Z]/K$. Then $J = (g_{ij,lk},\ f)$ where $g_{ij,lk} = (\{(x_{ij} - y_{ij})z_{lk} - (x_{lk} - y_{lk})z_{ij}\})$ and

$$f = \begin{vmatrix} x_{11} & x_{12} & x_{13} \\ z_{21} & z_{22} & z_{23} \\ y_{31} & y_{32} & y_{33} \end{vmatrix} + \begin{vmatrix} x_{11} & x_{12} & x_{13} \\ x_{21} & x_{22} & x_{23} \\ z_{31} & z_{32} & z_{33} \end{vmatrix}.$$

$K = (J,\ h)$ where

$$h = \begin{vmatrix} z_{11} & z_{12} & z_{13} \\ z_{21} & z_{22} & z_{23} \\ y_{31} & y_{32} & y_{33} \end{vmatrix} + \begin{vmatrix} z_{11} & z_{12} & z_{13} \\ y_{21} & y_{22} & y_{23} \\ z_{31} & z_{32} & z_{33} \end{vmatrix} + \begin{vmatrix} x_{11} & x_{12} & x_{13} \\ z_{21} & z_{22} & z_{23} \\ z_{31} & z_{32} & z_{33} \end{vmatrix}.$$

The remaining part of this section is devoted to prove Theorem 2.1. In the course of this we also describe the defining equations of $\mathcal{R}(\mathbb{D})$. We identify some specific equations in the defining ideal $\mathcal{J}$ of $\mathrm{Sym}(\mathbb{D})$. In order to clarify the notations, we define matrices that will be used repeatedly.

**Definition 2.3.** Let $X = [x_{ij}]$, $Y = [y_{ij}]$ , $1 \le i \le m, 1 \le j \le n$, be $m$ by $n$ matrices, and $X^{l,k}_{a_1 \dots a_s} = [x_{ia_i}]$, $Y^{l,k}_{a_1 \dots a_s} = [y_{ia_i}]$, $l \le i \le k$, $1 \le a_i \le n$, $X^{l,k}_{1 \dots \hat{s} \dots n} = [x_{ij}]$, $l \le i \le k$, $1 \le j \le n, j \ne s$ be submatrices. For the convenience of notation, we write $\det M = |M|$ when $M$ is a square matrix. We set determinate of a 0 by 0 matrix equal to 1. We also write

$$\begin{bmatrix} X^{1,j} \\ Y^{j+1,m} \end{bmatrix}_{a_1 \dots a_m} = \begin{bmatrix} x_{1a_1} & \cdots & x_{1a_m} \\ \vdots & & \vdots \\ x_{ja_1} & \cdots & x_{ja_m} \\ y_{j+1a_1} & \cdots & y_{j+1a_m} \\ \vdots & & \vdots \\ y_{ma_1} & \cdots & y_{ma_m} \end{bmatrix}$$

as a matrix with different variables.

The following is a well-known fact: writing a matrix with variable $y$'s as a matrix of variables $x$'s and a combination of differences of $x$'s and $y$'s.

**Lemma 2.4.** *Let $X$ and $Y$ be $n \times n$ matrices. With notation as above*

$$|Y| = |X| + \sum_{i=1}^{n} \sum_{j=1}^{n} (-1)^{i+j} \begin{vmatrix} Y^{1,i-1}_{1 \dots \hat{j} \dots n} \\ X^{i+1,n}_{1 \dots \hat{j} \dots n} \end{vmatrix} (y_{ij} - x_{ij})$$

*Proof.* We prove this by inducting on $n$. When $n = 1$, the claim is trivial. By induction, we have

$$|Y^{2,n}_{1 \dots \hat{k} \dots n}| = |X^{2,n}_{1 \dots \hat{k} \dots n}| + \sum_{i=2}^{n} \sum_{j=1}^{k-1} (-1)^{i+j+1} \begin{vmatrix} Y^{2,i-1}_{1 .. \hat{j} \dots . \hat{k} \dots . n} \\ X^{i+1,n}_{1 \dots \hat{j} \dots \hat{k} \dots n} \end{vmatrix} (y_{ij} - x_{ij})$$

$$+\sum_{i=2}^{n}\sum_{j=k+1}^{n}(-1)^{i+j}\begin{vmatrix} Y^{2,i-1}_{1..\hat{k}....\hat{j}....n} \\ X^{i+1,n}_{1...\hat{k}...\hat{j}...n} \end{vmatrix}(y_{ij}-x_{ij}).$$

Hence

$$\begin{aligned}
|Y| &= \textstyle\sum_{k=1}^{n}(-1)^{k+1}y_{1k}|Y^{2,n}_{1...\hat{k}...n}| \\
&= \textstyle\sum_{k=1}^{n}(-1)^{k+1}y_{1k}\left(|X^{2,n}_{1...\hat{k}...n}| + \displaystyle\sum_{i=2}^{n}\sum_{j=1}^{k-1}(-1)^{i+j+1}\begin{vmatrix} Y^{2,i-1}_{1..\hat{j}....\hat{k}....n} \\ X^{i+1,n}_{1...\hat{j}...\hat{k}...n} \end{vmatrix}(y_{ij}-x_{ij})\right. \\
&\qquad \left. + \textstyle\sum_{i=2}^{n}\sum_{j=k+1}^{n}(-1)^{i+j}\begin{vmatrix} Y^{2,i-1}_{1..\hat{k}....\hat{j}....n} \\ X^{i+1,n}_{1...\hat{k}...\hat{j}...n} \end{vmatrix}(y_{ij}-x_{ij})\right).
\end{aligned}$$

$$\begin{aligned}
&\overset{(1)}{=} \textstyle\sum_{k=1}^{n}(-1)^{k+1}x_{1k}|X^{2,n}_{1...\hat{k}...n}| + \sum_{k=1}^{n}(-1)^{k+1}(y_{1k}-x_{1k})|X^{2,n}_{1...\hat{k}...n}| \\
&\quad + \textstyle\sum_{k=1}^{n}(-1)^{k+1}y_{1k}\left(\sum_{i=2}^{n}\sum_{j=1}^{k-1}(-1)^{i+j+1}\begin{vmatrix} Y^{2,i-1}_{1..\hat{j}....\hat{k}....n} \\ X^{i+1,n}_{1...\hat{j}...\hat{k}...n} \end{vmatrix}(y_{ij}-x_{ij})\right. \\
&\qquad \left. + \textstyle\sum_{i=2}^{n}\sum_{j=k+1}^{n}(-1)^{i+j}\begin{vmatrix} Y^{2,i-1}_{1..\hat{k}....\hat{j}....n} \\ X^{i+1,n}_{1...\hat{k}...\hat{j}...n} \end{vmatrix}(y_{ij}-x_{ij})\right)
\end{aligned}$$

$$\begin{aligned}
&= |X| + \sum_{k=1}^{n}(-1)^{k+1}(y_{1k}-x_{1k})|X^{2,n}_{1...\hat{k}...n}| \\
&\quad + \sum_{k=1}^{n}(-1)^{k+1}y_{1k}\left(\sum_{i=2}^{n}\sum_{j=1}^{k-1}(-1)^{i+j+1}\begin{vmatrix} Y^{2,i-1}_{1..\hat{j}....\hat{k}....n} \\ X^{i+1,n}_{1...\hat{j}...\hat{k}...n} \end{vmatrix}(y_{ij}-x_{ij})\right. \\
&\quad \left. + \sum_{i=2}^{n}\sum_{j=k+1}^{n}(-1)^{i+j}\begin{vmatrix} Y^{2,i-1}_{1..\hat{k}....\hat{j}....n} \\ X^{i+1,n}_{1...\hat{k}...\hat{j}...n} \end{vmatrix}(y_{ij}-x_{ij})\right).
\end{aligned}$$

Here equality (1) comes from adding the extra term $\sum_{j=1}^{n}(-1)^{j+1}x_{1j}|X^{2,n}_{1...\hat{j}...n}|$ and then subtracting it from $\sum_{j=1}^{n}(-1)^{j+1}y_{1j}|X^{2,n}_{1...\hat{j}...n}|$. Notice that we have the third and forth sum as the following:

$$\begin{aligned}
&\sum_{k=1}^{n}(-1)^{k+1}y_{1k}\left(\textstyle\sum_{i=2}^{n}\sum_{j=1}^{k-1}(-1)^{i+j+1}\begin{vmatrix} Y^{2,i-1}_{1..\hat{j}....\hat{k}....n} \\ X^{i+1,n}_{1...\hat{j}...\hat{k}...n} \end{vmatrix}(y_{ij}-x_{ij})\right. \\
&\qquad \left. +\sum_{i=2}^{n}\sum_{j=k+1}^{n}(-1)^{i+j}\begin{vmatrix} Y^{2,i-1}_{1..\hat{k}....\hat{j}....n} \\ X^{i+1,n}_{1...\hat{k}...\hat{j}...n} \end{vmatrix}(y_{ij}-x_{ij})\right) \\
&= \sum_{i=2}^{n}\sum_{j=1}^{n}(-1)^{i+j}(y_{i,j}-x_{i,j})\left(\sum_{k=1}^{j-1}(-1)^{k+1}y_{1k}\begin{vmatrix} Y^{2,i-1}_{1...\hat{k}..\hat{j}...n} \\ X^{i+1,n}_{1...\hat{k}..\hat{j}...n} \end{vmatrix}\right.
\end{aligned}$$

$$
\begin{aligned}
&+\sum_{k=j+1}^{n}(-1)^{k}y_{1k}\begin{vmatrix} Y^{2,i-1}_{1\ldots\hat{j}..\hat{k}\ldots n} \\ X^{i+1,n}_{1\ldots\hat{j}..\hat{k}\ldots n} \end{vmatrix}\Bigg) \\
= &\sum_{i=2}^{n}\sum_{j=1}^{n}(-1)^{i+j}\begin{vmatrix} Y^{1,i-1}_{1\ldots\hat{j}\ldots n} \\ X^{i+1,n}_{1\ldots\hat{j}\ldots n} \end{vmatrix}(y_{ij}-x_{ij}).
\end{aligned}
$$

Now we can put everything together, we obtain

$$
\begin{aligned}
|Y| &= |X|+\sum_{j=1}^{n}(-1)^{j+1}|X^{2,n}_{1..\hat{j}\ldots n}|(y_{1j}-x_{1j}) \\
&\quad+\sum_{i=2}^{n}\sum_{j=1}^{n}(-1)^{i+j}\begin{vmatrix} Y^{1,i-1}_{1\ldots\hat{j}\ldots n} \\ X^{i+1,n}_{1\ldots\hat{j}\ldots n} \end{vmatrix}(y_{ij}-x_{ij}) \\
&= |X|+\sum_{i=1}^{n}\sum_{j=1}^{n}(-1)^{i+j}\begin{vmatrix} Y^{1,i-1}_{1\ldots\hat{j}\ldots n} \\ X^{j+1,n}_{1\ldots\hat{j}\ldots n} \end{vmatrix}(y_{ij}-x_{ij}).
\end{aligned}
$$

□

In the following lemma, we define those special equations that we consider and we show that those equations are in the defining ideal of symmetric algebra of $\mathbb{D}$.

**Lemma 2.5.** *Let $X_{a_1\ldots a_{s_1}}$ be the $s_1$ by $s_1$ submatrix of $X_{s_1t_1}$ with columns $a_1,\ldots,a_{s_1}$, $Y_{b_1\ldots b_{s_2}}$ the $s_2$ by $s_2$ submatrix of $Y_{s_2t_2}$ with columns $b_1,\ldots,b_{s_2}$, $X^{l,k}_{a_1\ldots a_{s_1}}$ the $k-l+1$ by $s_1$ submatrix of $X$ with rows $l$, $l+1,..,k$ and columns $a_1,\ldots,a_{s_1}$, and similarly for $Y$ and $Z$.*

*We define*

$$
g_{ij,lk}=\begin{vmatrix} z_{ij} & z_{lk} \\ x_{ij}-y_{ij} & x_{lk}-y_{lk} \end{vmatrix}
$$

$$
f_{a_1,\ldots,a_{s_1}}=\sum_{q=1}^{s_2}(-1)^{q+1}\left|\begin{bmatrix} Z^{q,q} \\ Y^{1,q-1} \\ X^{q+1,m} \end{bmatrix}_{a_1\ldots a_{s_1}}\right|,
$$

*where $1\le a_1<a_2<\ldots<a_{s_1}\le\min(t_1,t_2)$ and $1\le i\le m$, $1\le l\le m$, $1\le j\le n$, $1\le k\le n$.*

*We write $\mathcal{L}=(I_{s_1}(X_{s_1t_1}),\ I_{s_2}(Y_{s_2t_2}),\ g_{ij,lk},\ f_{a_1,\ldots,a_{s_1}})$, which is an ideal of $k[X_{mn},Y_{mn},Z_{mn}]$. Then $\mathcal{L}\subset\mathcal{J}$.*

*Proof.* We can see $|X_{a_1\ldots a_{s_1}}|$, $|Y_{b_1\ldots b_{s_2}}|$, $g_{ij,lk}$'s are in $\mathcal{J}$. Notice that when $t_2<s_1$, by the way we define $f_{a_1,\ldots,a_{s_1}}$, this is an empty condition. When $t_2\ge s_1$, we substitute $z_{ij}$ via $x_{ij}-y_{ij}$ and use Lemma 2.4, we can see $f$'s are in $\mathcal{J}$. □

The next Theorem immediate implies Theorem 2.1.

**Theorem 2.6.** *The ideal $\mathcal{L}$ is the defining ideal of $\mathcal{R}(\mathbb{D})$ and $\mathbb{D}$ is of linear type.*

The following lemma is the key step to prove the main theorem. The proof of the lemma is given in the Section 4. It involves finding a Gröbner basis of the ideal.

**Lemma 2.7.** *The variable $x_{11}$ is a non zero-divisor of the quotient ring $k[X,Y,Z]/\mathcal{L}$.*

*Proof of Theorem* 2.6: From Lemma 2.5, we have $\mathcal{L} \subset \mathcal{J} \subset \mathcal{K}$, where $\mathcal{J}$ is the defining ideal of $\mathrm{Sym}(\mathbb{D})$. We would like to show $\mathcal{L} = \mathcal{K}$ and as a consequence, $\mathcal{L} = \mathcal{J} = \mathcal{K}$, i.e. $\mathbb{D}$ is an ideal of linear type. By Lemma 2.7, $x_{11}$ is a non zero-divisor on $k[X,Y,Z]/\mathcal{L}$. Changing the roles of $X$ and $Y$, we also obtain that $y_{11}$ is a non-zero divisor on $k[X,Y,Z]/\mathcal{L}$. Since $\mathcal{K}$ is a prime ideal, it suffices to show that $\mathcal{L}_{x_{11}y_{11}} = \mathcal{K}_{x_{11}y_{11}}$. The latter holds by induction on the size of the matrix $X$.

To explain this, we consider the $(m-1)\times(n-1)$ matrices of variables $X' = [x'_{ij}]$ , $Y' = [y_{ij}]$, $Z' = [z_{ij}]$, $2 \le i \le m$, $2 \le j \le n$. We define a natural isomorphism $\phi$ from $k[\{x_{ij}\}_{i=1 \text{ or } j=1}, X']_{x_{11}}$ to $k[X]_{x_{11}}$ via $\phi(x_{ij}) = x_{ij}$ when $i=1$ or $j=1$, and $\phi(x'_{ij}) = x_{ij} - x_{i1}x_{1j}/x_{11}$ when $i \ne 1$ and $j \ne 1$. Let

$$R'_1 = k[\{x_{ij}\}_{i=1 \text{ or } j=1}, X']_{x_{11}}/I' \cong R_{1x_{11}},$$

$$R'_2 = k[\{x_{ij}\}_{i=1 \text{ or } j=1}, X']_{x_{11}}/J' \cong R_{2x_{11}},$$

where $I' = I_{s_1-1}(X'_{s_1-1,t_1-1})$, $J' = I_{s_2-1}(X'_{s_2-1,t_2-1})$ and

$$\begin{aligned} S' &:= k[\{x_{ij}\}_{i=1 \text{ or } j=1}, X', \{y_{ij}\}_{i=1 \text{ or } j=1}, Y']_{x_{11}y_{11}}/(I',J') \\ &\cong R'_1 \otimes R'_2 \cong R_{1x_{11}} \otimes R_{2x_{11}}. \end{aligned}$$

Then we have

$$\begin{aligned} \widetilde{\mathbb{D}}' &= (\{x_{ij} - y_{ij}\}_{i=1 \text{ or } j=1}, \{x'_{ij} - y'_{ij}\}_{2\le i\le m,\ 2\le j\le n}) \\ &\cong \mathbb{D} = (\{x_{ij} - y_{ij}\}_{1\le i\le m,\ 1\le j\le n}), \end{aligned}$$

and $T' = S'[\{z_{ij}\}_{i=1 \text{ or } j=1}, Z'] \cong T_{x_{11}y_{11}}$ by the map $\overline{\phi}$ defined by $\overline{\phi}(x_{ij}) = x_{ij}$, $\overline{\phi}(y_{ij}) = y_{ij}$ and $\overline{\phi}(z_{ij}) = z_{ij}$ when $i=1$ or $j=1$, and $\overline{\phi}(x'_{ij}) = x_{ij} - x_{i1}x_{1j}/x_{11}$, $\overline{\phi}(y'_{ij}) = y_{ij} - y_{i1}y_{1j}/y_{11}$, and
$\overline{\phi}(z'_{ij}) = z_{ij} - y_{i1}z_{1j}/y_{11} - x_{1j}z_{i1}/y_{11} + x_{i1}x_{1j}z_{11}/x_{11}y_{11}$ when $i \ne 1$ and $j \ne 1$. Let $\phi'$ denote the induced map of $\overline{\phi}$ from $\mathcal{R}_{S'}(\widetilde{\mathbb{D}}')$ to $\mathcal{R}_{S_{x_{11}y_{11}}}(\mathbb{D})$. Let $\psi$ and $\psi'$ denote the map from $T_{x_{11}y_{11}}$to $\mathcal{R}_{S_{x_{11}y_{11}}}(\mathbb{D})$ and $T'$ to $\mathcal{R}_{S'}(\widetilde{\mathbb{D}}')$. We obtain the following diagram

$$\begin{tikzcd} T' \arrow[r, "\overline{\phi}"] \arrow[d, "\psi'"'] & T_{x_{11}y_{11}} \arrow[d, "\psi"] \\ \mathcal{R}_{S'}(\mathbb{D}') \arrow[r, "\phi'"'] & \mathcal{R}_{S_{x_{11}y_{11}}}(\mathbb{D}) \end{tikzcd}$$

To show the diagram commute, it is enough to show $\phi'(\psi'(z'_{ij})) = \psi(\overline{\phi}(z'_{ij}))$. Since

$$\begin{aligned} \psi(\overline{\phi}(z'_{ij})) &= \psi(z_{ij} - y_{i1}z_{1j}/y_{11} - x_{1j}z_{i1}/y_{11} + x_{i1}x_{1j}z_{11}/x_{11}y_{11}) \\ &= x_{ij} - y_{ij} - y_{i1}(x_{1j} - y_{1j})/y_{11} - x_{1j}(x_{i1} - y_{i1})/y_{11} \\ &\quad +x_{i1}x_{1j}(x_{11} - y_{11})/x_{11}y_{11}, \end{aligned}$$

and

$$\begin{aligned} \phi'(\psi'(z'_{ij})) &= \phi'(x'_{ij} - y'_{ij}) \\ &= x_{ij} - y_{ij} - x_{i1}x_{1j}/x_{11} + y_{i1}y_{1j}/y_{11} \\ &= x_{ij} - y_{ij} - y_{i1}(x_{1j} - y_{1j})/y_{11} - x_{1j}(x_{i1} - y_{i1})/y_{11} \\ &\quad +x_{i1}x_{1j}(x_{11} - y_{11})/x_{11}y_{11}. \end{aligned}$$

Hence $\phi'$ is an isomorphism. Let $\mathbb{D}' = (\{x'_{ij} - y'_{ij}\}_{2\leq i\leq m,\ 2\leq j\leq n})$ then by the induction hypothesis, the defining ideal of $\mathcal{R}_{S'}(\mathbb{D}')$ in $T'$ is of the form $\mathcal{L}' = \{I_{s_1-1}(X'_{s_1-1,t_1-1}),\ I_{s_2-1}(Y'_{s_2-1,t_2-1}), g'_{ij,lk}, f'_{a_2,\ldots,a_{s-1}}\}$, where

$$g'_{ij,lk} = \begin{vmatrix} z'_{ij} & z'_{lk} \\ x'_{ij} - y'_{ij} & x'_{lk} - y'_{lk} \end{vmatrix}$$

$$f'_{a_2,\ldots,a_{s_1}} = \sum_{q=2}^{s_2} (-1)^{q+1} \left| \begin{bmatrix} Z'^{q,q} \\ Y'^{2,q-1} \\ X'^{q+1,m} \end{bmatrix}_{a_2\ldots a_{s_1}} \right|$$

with $2 \leq a_1 < a_2 < \ldots < a_{s_1} \leq \min\{t_1, t_2\}$ and $2 \leq i \leq m$, $2 \leq l \leq m$, $2 \leq j \leq n$, $2 \leq k \leq n$. Let $W$ denote the set of Koszul relations:

$${g^1}_{ij,lk} = \begin{vmatrix} z_{ij} & z'_{lk} \\ x_{ij} - y_{ij} & x'_{lk} - y'_{lk} \end{vmatrix}$$

with $i = 1$ or $j = 1$ and

$${g^2}_{ij,lk} = \begin{vmatrix} z_{ij} & z_{lk} \\ x_{ij} - y_{ij} & x_{lk} - y_{lk} \end{vmatrix}$$

with $i = 1$ or $j = 1$ and $l = 1$ or $k = 1$. Then $(\mathcal{L}', W)$ is the defining ideal of $\mathcal{R}_{S'}(\widetilde{\mathbb{D}'}) \cong \mathcal{R}_{S_{x_{11}y_{11}}}(\mathbb{D})$. Once we show that $\overline{\phi}(\mathcal{L}', W) \subset \mathcal{L}_{x_{11}y_{11}}$, then $\mathcal{L}_{x_{11}y_{11}} = \mathcal{K}_{x_{11}y_{11}}$.

From the way we define the map $\overline{\phi}$, we have

$$\overline{\phi}(I_{s_1-1}(X'_{s_1-1,t_1-1}),\ I_{s_2-1}(Y'_{s_2-1,t_2-1}), g'_{ij,lk}, W) \subset \mathcal{L}_{x_{11}y_{11}}.$$

Notice the following equality:

$$\overline{\phi}(f'_{a_2,\dots,a_{s_1}}) \quad = \quad \frac{1}{y_{11}} f_{1,a_2,\dots,a_{s_1}} - \frac{z_{11}}{x_{11}y_{11}} |X_{1a_2\dots a_{s_1}}|,$$

hence $\overline{\phi}(f'_{a_2,\dots,a_{s_1}}) \in \mathcal{L}_{x_{11}y_{11}}$.

$\square$

## 3. SOME LINEAR ALGEBRA

This section is a reminder of some linear algebra properties. We will use those properties in the proof of Section 4.

The following lemma writes the determinant of a certain in $x$ and $y$ variables in term of $y$ variables and differences $x_{ij} - y_{ij}$.

**Lemma 3.1.** *With notation as 2.3,*

$$\begin{vmatrix} & & Y^{1,i-1}_{1,\dots,n} & & & \\ y_{i,1} & \cdots & y_{i,j} & x_{i,j+1} & \cdots & x_{i,n} \\ & & X^{i+1,n}_{1,\dots,n} & & & \end{vmatrix}$$

$$\begin{aligned} = \quad & |Y| + \sum_{k=j+1}^{n} (-1)^{i+k} \begin{vmatrix} Y^{1,i-1}_{1..\hat{k}..n} \\ X^{i+1,n}_{1\dots\hat{k}\dots n} \end{vmatrix} (x_{ik} - y_{ik}) \\ & + \sum_{l=i+1}^{n} \sum_{k=1}^{n} (-1)^{l+k} \begin{vmatrix} Y^{1,l-1}_{1\dots\hat{k}\dots n} \\ X^{l+1,n}_{1\dots\hat{k}\dots n} \end{vmatrix} (x_{lk} - y_{lk}). \end{aligned}$$

*Proof.* We will show this by inducting on $i$. When $i = 1$, we have

$$\begin{vmatrix} y_{1,1} \dots y_{1,j} x_{1,j+1} .. x_{1,n} \\ X^{2,n} \end{vmatrix}$$

$$\begin{aligned} = \quad & \sum_{k=1}^{j} (-1)^{k+1} y_{1,k} |X^{2,n}_{1\dots\hat{k}\dots n}| + \sum_{k=j+1}^{n} (-1)^{k+1} x_{1,k} |X^{2,n}_{1\dots\hat{k}\dots n}| \\ = \quad & \sum_{k=1}^{j} (-1)^{k+1} (y_{1,k} - x_{1,k}) |X^{2,n}_{1\dots\hat{k}\dots n}| + \sum_{k=1}^{n} (-1)^{k+1} x_{1,k} |X^{2,n}_{1\dots\hat{k}\dots n}| \\ = \quad & \sum_{k=1}^{j} (-1)^{k+1} (y_{1,k} - x_{1,k}) |X^{2,n}_{1\dots\hat{k}\dots n}| + |X| \quad . \end{aligned}$$

Now Lemma 2.4 gives

$$|X| = |Y| + \sum_{l=1}^{n} \sum_{j=1}^{n} (-1)^{l+j} \begin{vmatrix} Y^{1,l-1}_{1\dots\hat{j}\dots n} \\ X^{l+1,n}_{1\dots\hat{j}\dots n} \end{vmatrix} (x_{lj} - y_{lj}).$$

Hence we have

$$\begin{aligned} &\begin{vmatrix} y_{1,1}\dots y_{1,j}x_{1,j+1}..x_{1,n} \\ X^{2,n} \end{vmatrix} \\ = &\sum_{k=1}^{j}(-1)^{k+1}(y_{1k}-x_{1k})|X^{2,n}_{1\dots\hat{k}\dots n}| + |Y| + \sum_{l=1}^{n}\sum_{j=1}^{n}(-1)^{l+j}\begin{vmatrix} Y^{1,l-1}_{1\dots\hat{j}\dots n} \\ X^{l+1,n}_{1\dots\hat{j}\dots n}\end{vmatrix}(x_{lj}-y_{lj}). \end{aligned}$$

Notice that

$$\begin{aligned} &\sum_{k=1}^{j}(-1)^{k+1}(y_{1k}-x_{1k})|X^{2,n}_{1\dots\hat{k}\dots n}| + \sum_{l=1}^{n}\sum_{j=1}^{n}(-1)^{l+j}\begin{vmatrix} Y^{1,l-1}_{1\dots\hat{j}\dots n} \\ X^{l+1,n}_{1\dots\hat{j}\dots n}\end{vmatrix}(x_{lj}-y_{lj}) \\ = &\sum_{k=1}^{j}(-1)^{k+1}(y_{1k}-x_{1k})|X^{2,n}_{1\dots\hat{k}\dots n}| + \sum_{j=1}^{n}(-1)^{j+1}|X^{2,n}_{1\dots\hat{j}\dots n}|(x_{1j}-y_{1j}) \\ &+\sum_{l=2}^{n}\sum_{j=1}^{n}(-1)^{l+j}\begin{vmatrix} Y^{1,l-1}_{1\dots\hat{j}\dots n} \\ X^{l+1,n}_{1\dots\hat{j}\dots n}\end{vmatrix}(x_{lj}-y_{lj}) \\ = &\sum_{k=j+1}^{n}(-1)^{k+1}(x_{1k}-y_{1k})|X^{2,n}_{1\dots\hat{k}\dots n}| + \sum_{l=2}^{n}\sum_{j=1}^{n}(-1)^{l+j}\begin{vmatrix} Y^{1,l-1}_{1\dots\hat{j}\dots n} \\ X^{l+1,n}_{1\dots\hat{j}\dots n}\end{vmatrix}(x_{lj}-y_{lj}) \quad . \end{aligned}$$

Hence

$$\begin{aligned} \begin{vmatrix} y_{1,1}\dots y_{1,j}x_{1,j+1}..x_{1,n} \\ X^{2,n} \end{vmatrix} = &|Y| + \sum_{k=j+1}^{n}(-1)^{k+1}(x_{1,k}-y_{1,k})|X^{2,n}_{1\dots\hat{k}\dots n}| \\ &+\sum_{l=2}^{n}\sum_{j=1}^{n}(-1)^{l+j}\begin{vmatrix} Y^{1,l-1}_{1\dots\hat{j}\dots n} \\ X^{l+1,n}_{1\dots\hat{j}\dots n}\end{vmatrix}(x_{lj}-y_{lj}). \end{aligned}$$

Now for the induction step, we assume

$$\begin{vmatrix} Y^{1,i-1} \\ X^{i,n} \end{vmatrix} = |Y| + \sum_{l=i}^{n}\sum_{j=1}^{n}(-1)^{l+j}\begin{vmatrix} Y^{1,l-1}_{1\dots\hat{j}\dots n} \\ X^{l+1,n}_{1\dots\hat{j}\dots n}\end{vmatrix}(x_{lj}-y_{lj}).$$

Therefore

$$\begin{aligned} &\begin{vmatrix} Y^{1,i-1} \\ y_{i,1}\dots y_{i,j}x_{i,j+1}..x_{i,n} \\ X^{i+1,n} \end{vmatrix} \\ = &\sum_{k=1}^{j}(-1)^{i+k}y_{i,k}\begin{vmatrix} Y^{1,i-1}_{1\dots\hat{k}\dots n} \\ X^{i+1,n}_{1\dots\hat{k}\dots n}\end{vmatrix} + \sum_{k=j+1}^{n}(-1)^{i+k}x_{i,k}\begin{vmatrix} Y^{1,i-1}_{1\dots\hat{k}\dots n} \\ X^{i+1,n}_{1\dots\hat{k}\dots n}\end{vmatrix} \\ = &\sum_{k=1}^{j}(-1)^{i+k}(y_{i,k}-x_{i,k})\begin{vmatrix} Y^{1,i-1}_{1\dots\hat{k}\dots n} \\ X^{i+1,n}_{1\dots\hat{k}\dots n}\end{vmatrix} + \sum_{k=1}^{n}(-1)^{i+k}x_{i,k}\begin{vmatrix} Y^{1,i-1}_{1\dots\hat{k}\dots n} \\ X^{i+1,n}_{1\dots\hat{k}\dots n}\end{vmatrix} \end{aligned}$$

$$
\begin{aligned}
&= \sum_{k=1}^{j}(-1)^{i+k}(y_{i,k}-x_{i,k}) \begin{vmatrix} Y^{1,i-1}_{1\ldots\hat{k}\ldots n} \\ X^{i+1,n}_{1\ldots\hat{k}\ldots n} \end{vmatrix} + \begin{vmatrix} Y^{1,i-1} \\ X^{i,n} \end{vmatrix} \\
&= \sum_{k=1}^{j}(-1)^{i+k}(y_{i,k}-x_{i,k}) \begin{vmatrix} Y^{1,i-1}_{1\ldots\hat{k}\ldots n} \\ X^{i+1,n}_{1\ldots\hat{k}\ldots n} \end{vmatrix} + |Y| \\
&\quad + \sum_{l=i}^{n}\sum_{j=1}^{n}(-1)^{l+j} \begin{vmatrix} Y^{1,l-1}_{1\ldots\hat{j}\ldots n} \\ X^{l+1,n}_{1\ldots\hat{j}\ldots n} \end{vmatrix} (x_{lj}-y_{lj}) \\
&= |Y| + \sum_{k=j+1}^{n}(-1)^{i+k}(x_{i,k}-y_{i,k}) \begin{vmatrix} Y^{1,i-1}_{1\ldots\hat{k}\ldots n} \\ X^{i+1,n}_{1\ldots\hat{k}\ldots n} \end{vmatrix} \\
&\quad + \sum_{l=i+1}^{n}\sum_{j=1}^{n}(-1)^{l+j} \begin{vmatrix} Y^{1,l-1}_{1\ldots\hat{j}\ldots n} \\ X^{l+1,n}_{1\ldots\hat{j}\ldots n} \end{vmatrix} (x_{lj}-y_{lj}).
\end{aligned}
$$

□

By the following the two lemmas, the $S$-pairs between the elements of the ideal $\mathcal{L}$ can be reduced using the Koszul relations.

**Lemma 3.2.** *Let* $1 \leq i,\ l \leq m,\ 1 \leq j,\ k \leq n,\ a_1 < a_2 < a_3$. *Let*

$$
g_{ij,lk} = \begin{vmatrix} z_{ij} & z_{lk} \\ x_{ij}-y_{ij} & x_{lk}-y_{lk} \end{vmatrix},\ M = \begin{vmatrix} z_{1a_1} & z_{1a_2} & z_{1a_3} \\ x_{1a_1} & x_{1a_2} & x_{1a_3} \\ y_{1a_1} & y_{1a_2} & y_{1a_3} \end{vmatrix}.
$$

*Then*

$$
M = y_{1a_1}g_{1a_2,1a_3} - y_{1a_2}g_{1a_1,1a_3} + y_{1a_3}g_{1a_1,1a_2}.
$$

*Proof.* One has

$$
M = \begin{vmatrix} z_{1a_1} & z_{1a_2} & z_{1a_3} \\ x_{1a_1}-y_{1a_1} & x_{1a_2}-y_{1a_2} & x_{1a_3}-y_{1a_3} \\ y_{1a_1} & y_{1a_2} & y_{1a_3} \end{vmatrix}
$$

$$
\begin{aligned}
&= y_{1a_1}\begin{vmatrix} z_{1a_2} & z_{1a_3} \\ x_{1a_2}-y_{1a_2} & x_{1a_3}-y_{1a_3} \end{vmatrix} - y_{1a_2}\begin{vmatrix} z_{1a_1} & z_{1a_3} \\ x_{1a_1}-y_{a_1} & x_{1a_3}-y_{1a_3} \end{vmatrix} \\
&\quad + y_{1a_3}\begin{vmatrix} z_{1a_1} & z_{1a_2} \\ x_{1a_1}-y_{1a_1} & x_{1a_2}-y_{1a_2} \end{vmatrix}
\end{aligned}
$$

□

**Lemma 3.3.** *Let* $g_{ij,lk}$ *defined as Lemma 3.2. Then*

$$\begin{aligned} M &= \begin{vmatrix} z_{1a_1} & z_{1a_2} \\ x_{2a_1} - y_{2a_1} & x_{2a_2} - y_{2a_2} \end{vmatrix} \\ &= g_{1a_1,2a_2} - g_{1a_2,2a_1} + \begin{vmatrix} x_{1a_1} - y_{1a_1} & x_{1a_2} - y_{1a_2} \\ z_{2a_1} & z_{2a_2} \end{vmatrix}. \end{aligned}$$

*Proof.* We have

$$\begin{aligned} M &= z_{1a_1}(x_{2a_2} - y_{2a_2}) - z_{1a_2}(x_{2a_1} - y_{2a_1}) \\ &= g_{1a_1,2a_2} + z_{2a_2}(x_{1a_1} - y_{1a_1}) - g_{1a_2,2a_1} - z_{2a_1}(x_{1a_2} - y_{1a_2}) \\ &= g_{1a_1,2a_2} - g_{1a_2,2a_1} + \left| \begin{bmatrix} x_{1a_1} - y_{1a_1} & x_{1a_2} - y_{1a_2} \\ z_{2a_1} & z_{2a_2} \end{bmatrix} \right|. \end{aligned}$$

□

## 4. Gröbner basis

This section is devoted to prove Lemma 2.7. We outline this section here. We will recall Buchberger's Criterion and give several lemmas that will help us reduce the computation of $S$-pairs between elements of $\mathcal{L}$. We define several equations and show those equations sit inside the ideal $\mathcal{L}$. Theorem 4.25 will give a Gröbner basis of $\mathcal{L}$ via a particular ordering. Actually, all the equations defined before Theorem 4.25 are all of elements of the Gröbner basis. The proof of Theorem 4.25 will be broken down as several lemmas computing the $S$-pairs of the elements and showing all of the reminders of $S$-pairs are zero.

Let $I = (g_1, ..., g_s)$ be an ideal in a polynomial ring. We define

$$\text{in}(g_j)/\text{GCD}(\text{in}(g_i), \text{in}(g_j)) = m_{ji},$$

$$\text{in}(g_i)/\text{GCD}(\text{in}(g_i), \text{in}(g_j)) = m_{ij},$$

and

$$m_{ji}g_i - m_{ij}g_j = \sum f_u^{(ij)} g_u + h_{g_ig_j}$$

where $\text{in}(m_{ji}g_i) > \text{in}(f_u^{(ij)}g_u)$ for all $u$.

**Theorem 4.1.** (Buchberger′s Criterion). *The elements* $g_1, ..., g_s$ *form a Gröbner basis if and only if* $h_{g_ig_j} = 0$ *for all* $i$ *and* $j$.

The polynomial $m_{ji}g_i - m_{ij}g_j$ is commonly referred to as the $S$-pair between $g_i$ and $g_j$ and $h_{g_ig_j}$ is called the remainder.

By using Buchberger's Criterion, we obtain several lemmas that will help in the computation of a Gröbner basis of $\mathcal{L}$. Sine we focus on the determinantal rings, the computation of $S$-pair between elements are involving the values of matrix determinate. For the computation purpose, we define the following definition.

**Definition 4.2.** Given two square free monomials $p_1$ and $p_2$ in $k[X]$ where $X$ is the $m$ by $n$ matrix of variables, we define $m_{12} = p_1/\mathrm{GCD}(p_1, p_2)$ and $m_{21} = p_2/\mathrm{GCD}(p_1, p_2)$. Assume $m_{12} = x_{u_1a_1}...x_{u_ra_r}$, $m_{21} = x_{v_1b_1}...x_{v_wb_w}$, then define the matrix

$$M_{12} := \begin{bmatrix} x_{u_1a_1} & \cdots & x_{u_1a_r} \\ x_{u_2a_1} & \cdots & x_{u_2a_r} \\ \vdots & & \vdots \\ x_{u_ra_1} & & x_{u_ra_r} \end{bmatrix}$$

and the matrix

$$M_{21} := \begin{bmatrix} x_{v_1b_1} & \cdots & x_{v_1b_w} \\ x_{v_2b_1} & \cdots & x_{v_2b_w} \\ \vdots & & \vdots \\ x_{v_wb_1} & & x_{v_wb_w} \end{bmatrix} \quad .$$

The following lemma helps us replace a polynomial with a leading term involving $x_{i,j}$'s by a polynomial with a leading term without involving $x_{ij}$'s.

**Lemma 4.3.** *Let $a_{s_1} < ... < a_1$, $1 \le r \le s_1$, and let $g_{i_1j_1,i_2j_2}$ be as defined in Lemma 2.5. Then*

$$\left| \begin{bmatrix} Y^{1,r-1} \\ Z^{r,r} \\ X^{r+1,s_1} \end{bmatrix}_{a_{s_1},\dots,a_1} \right| = \left| \begin{bmatrix} Y^{1,r-1} \\ Z^{r,r} \\ Y^{r+1,s_1} \end{bmatrix}_{a_{s_1},\dots,a_1} \right| + \sum_{u=r+1}^{s_1} \left| \begin{bmatrix} Y^{1,r-1} \\ X^{r,r} - Y^{r,r} \\ Y^{r+1,u-1} \\ Z^{u,u} \\ X^{u+1,s_1} \end{bmatrix}_{a_{s_1},\dots,a_1} \right|$$

$$+ \sum_{u=r+1}^{s_1} \sum_{\{c_1,c_2,d_1,\dots,d_{s_1-2}\}=\{a_1,\dots,a_{s_1}\}} \pm \left(g_{rc_1,uc_2} - g_{rc_2,uc_1}\right) \left| \begin{bmatrix} Y^{1,r-1} \\ Y^{r+1,u-1} \\ X^{u+1,s_1} \end{bmatrix}_{d_1,\dots,d_{s_1-2}} \right| .$$

*Proof.* For the purpose of this proof we drop the column indices. We use Lemma 3.1 to obtain

$$\left| \begin{bmatrix} Y^{1,r-1} \\ Z^{r,r} \\ X^{r+1,s_1} \end{bmatrix} \right| = \left| \begin{bmatrix} Y^{1,r-1} \\ Z^{r,r} \\ Y^{r+1,s_1} \end{bmatrix} \right| + \sum_{u=r+1}^{s_1} \left| \begin{bmatrix} Y^{1,r-1} \\ Z^{r,r} \\ Y^{r+1,u-1} \\ X^{u,u} - Y^{u,u} \\ X^{u+1,s_1} \end{bmatrix} \right| .$$

Notice that

$$
\begin{aligned}
&\left|\begin{bmatrix} Y^{1,r-1} \\ Z^{r,r} \\ Y^{r+1,u-1} \\ X^{u,u}-Y^{u,u} \\ X^{u+1,s_1} \end{bmatrix}\right| \\
= & \sum_{\{c_1,c_2,d_1,\dots,d_{s_1-2}\}=\{a_1,\dots,a_{s_1}\}} \\
& \pm\left|\begin{bmatrix} z_{rc_1} & z_{rc_2} \\ x_{uc_1}-y_{uc_1} & x_{uc_2}-y_{uc_2} \end{bmatrix}\right| \left|\begin{bmatrix} Y^{1,r-1} \\ Y^{r+1,u-1} \\ X^{u+1,s_1} \end{bmatrix}_{d_1,\dots,d_{s_1-2}}\right| \\
= & \sum_{\{c_1,c_2,d_1,\dots,d_{s_1-2}\}=\{a_1,\dots,a_{s_1}\}} \pm\Bigg(g_{rc_1,uc_2}-g_{rc_2,uc_1} \\
& +\left|\begin{bmatrix} x_{rc_1}-y_{rc_1} & x_{rc_2}-y_{rc_2} \\ z_{uc_1} & z_{uc_2} \end{bmatrix}\right|\Bigg) \left|\begin{bmatrix} Y^{1,r-1} \\ Y^{r+1,u-1} \\ X^{u+1,s_1} \end{bmatrix}_{d_1,\dots,d_{s_1-2}}\right| \\
= & \sum_{u=r+1}^{s_1}\sum_{\{c_1,c_2,d_1,\dots,d_{s_1-2}\}=\{a_1,\dots,a_{s_1}\}} \\
& \pm(g_{rc_1,uc_2}-g_{rc_2,uc_1}) \left|\begin{bmatrix} Y^{1,r-1} \\ Y^{r+1,u-1} \\ X^{u+1,s_1} \end{bmatrix}_{d_1,\dots,d_{s_1-2}}\right| \\
& +\textstyle\sum_{u=r+1}^{s_1} \left|\begin{bmatrix} Y^{1,r-1} \\ X^{r,r}-Y^{r,r} \\ Y^{r+1,u-1} \\ Z^{u,u} \\ X^{u+1,s_1} \end{bmatrix}\right| .
\end{aligned}
$$

□

The determinant in the following lemma appears in many cases in the computing of Gröbner basis. This lemma enables the determinant to be written as a combination of elements of $I_{s_2}(Y)$ and $g_{i_1j_1,i_2j_2}$.

**Lemma 4.4.** *Let* $a_1 < ... < a_{s_1+1}$, *and* $1 \leq r \leq s_1$. *One has*

$$\sum_{u=r}^{s_2} \left| \begin{bmatrix} X^{r,r} \\ Y^{1,u-1} \\ Z^{u,u} \\ X^{u+1,s_1} \end{bmatrix}_{a_1,\ldots,a_{s_1+1}} \right| \in I_{s_2}(Y) + (g_{i_1 j_1, i_2 j_2} \,|\, 1 \leq i_v \leq m,\, 1 \leq j_v \leq n, v = 1, 2).$$

*Proof.* The column indices are omitted again. First we write

$$\sum_{u=r}^{s_2} \left| \begin{bmatrix} X^{r,r} \\ Y^{1,u-1} \\ Z^{u,u} \\ X^{u+1,s_1} \end{bmatrix} \right| = \left| \begin{bmatrix} X^{r,r} \\ Y^{1,r-1} \\ Z^{r,r} \\ X^{r+1,s_1} \end{bmatrix} \right| + \sum_{u=r+1}^{s_2} \left| \begin{bmatrix} X^{r,r} \\ Y^{1,u-1} \\ Z^{u,u} \\ X^{u+1,s_1} \end{bmatrix} \right|$$

then using Lemma 3.2 and 4.3, we obtain

$$\left| \begin{bmatrix} X^{r,r} \\ Y^{1,r-1} \\ Z^{r,r} \\ X^{r+1,s_1} \end{bmatrix} \right| = \left| \begin{bmatrix} X^{r,r} - Y^{r,r} \\ Y^{1,r-1} \\ Z^{r,r} \\ X^{r+1,s_1} \end{bmatrix} \right| + \left| \begin{bmatrix} Y^{r,r} \\ Y^{1,r-1} \\ Z^{r,r} \\ X^{r+1,s_1} \end{bmatrix} \right|$$

$$= \sum_{\{c_1,c_2,d_1,\ldots,d_{s_1-1}\}=\{a_1,\ldots,a_{s_1+1}\}} \pm\, g_{rc_1,rc_2} \left| \begin{bmatrix} Y^{1,r-1} \\ X^{r+1,s_1} \end{bmatrix}_{c_1,\ldots,c_{s_1-1}} \right| + \left| \begin{bmatrix} Y^{r,r} \\ Y^{1,r-1} \\ Z^{r,r} \\ Y^{r+1,s_1} \end{bmatrix} \right|$$

$$+ \sum_{u=r+1}^{s_2} \sum_{\{c_1,c_2,d_1,\ldots,d_{s_1-1}\}=\{a_1,\ldots,a_{s_1+1}\}} \pm (g_{rc_1,uc_2} - g_{rc_2,uc_1}) \left| \begin{bmatrix} Y^{r,r} \\ Y^{1,r-1} \\ Y^{r+1,u-1} \\ X^{u+1,s_1} \end{bmatrix}_{d_1,\ldots,d_{s_1-1}} \right|$$

$$+ \sum_{u=r+1}^{s_2} \left| \begin{bmatrix} Y^{r,r} \\ Y^{1,r-1} \\ X^{r,r} - Y^{r,r} \\ Y^{r+1,u-1} \\ Z^{u,u} \\ X^{u+1,s_1} \end{bmatrix} \right| + \sum_{u=s_2+1}^{s_1} \left| \begin{bmatrix} Y^{r,r} \\ Z^{r} \\ Y^{1,r-1} \\ Y^{r+1,u-1} \\ X^{u,u} - Y^{u,u} \\ X^{u+1,s_1} \end{bmatrix} \right|.$$

$\alpha$ is defined as given :

$$\alpha = \sum_{\{c_1,c_2,d_1,\ldots,d_{s_1-1}\}=\{a_1,\ldots,a_{s_1+1}\}} \pm g_{rc_1,rc_2} \left| \begin{bmatrix} Y^{1,r-1} \\ X^{r+1,s_1} \end{bmatrix}_{c_1,\ldots,c_{s_1-1}} \right| + \left| \begin{bmatrix} Y^{r,r} \\ Y^{1,r-1} \\ Z^{r,r} \\ Y^{r+1,s_1} \end{bmatrix} \right|$$

$$+\sum_{u=r+1}^{s_2}\sum_{\{c_1,c_2,d_1,\ldots,d_{s_1-1}\}=\{a_1,\ldots,a_{s_1+1}\}}\pm(g_{rc_1,uc_2}-g_{rc_2,uc_1})\left|\begin{bmatrix} Y^{r,r}\\ Y^{1,r-1}\\ Y^{r+1,u-1}\\ X^{u+1,s_1}\end{bmatrix}_{d_1,\ldots,d_{s_1-1}}\right|$$

$$+\textstyle\sum_{u=s_2+1}^{s_1}\left|\begin{bmatrix} Y^{r,r}\\ Z^{r}\\ Y^{1,r-1}\\ Y^{r+1,u-1}\\ X^{u,u}-Y^{u,u}\\ X^{u+1,s_1}\end{bmatrix}\right|.$$

This shows the element $\alpha$ is in $I_{s_2}(Y_{s_2t_2})+(g_{i_1j_1,i_2j_2})$. After removing the repeated row $y_r$, we have

$$\sum_{u=r+1}^{s_2}\left|\begin{bmatrix} Y^{r,r}\\ Y^{1,r-1}\\ X^{r,r}-Y^{r,r}\\ Y^{r+1,u-1}\\ Z^{u,u}\\ X^{u+1,s_1}\end{bmatrix}\right| = \sum_{u=r+1}^{s_2}\left|\begin{bmatrix} Y^{r,r}\\ Y^{1,r-1}\\ X^{r,r}\\ Y^{r+1,u-1}\\ Z^{u,u}\\ X^{u+1,s_1}\end{bmatrix}\right|$$

$$= \quad -\sum_{u=r+1}^{s_2}\left|\begin{bmatrix} X^{r,r}\\ Y^{1,r-1}\\ Y^{r,r}\\ Y^{r+1,u-1}\\ Z^{u,u}\\ X^{u+1,s_1}\end{bmatrix}\right| = -\sum_{u=r+1}^{s_2}\left|\begin{bmatrix} X^{r,r}\\ Y^{1,u-1}\\ Z^{u,u}\\ X^{u+1,s_1}\end{bmatrix}\right|.$$

□

In order to simplify the notation and the computation, we define notation to keep track of sums of determinants.

**Definition 4.5.** Let $G$ be a collection of polynomials in the ring $k[X,Y,Z]$ with $X$, $Y$, $Z$ as $m$ by $n$ matrices of variables over the field $k$. Let $\{P^u_{a_1,\ldots,a_{q_u}}\}_{u\in I}$ be an element of $G$ such that each $P^u_{a_1,\ldots,a_{q_u}}$ is the sum of determinants $P^u_i$ of $m$ by $n$ matrices with the same column indices, $a_1,\ldots,a_{q_u}$, in variables $X$, $Y$ and $Z$. Denote $P^u_{a_1,\ldots,a_{q_u}}=\sum_{i=1}^{p_u}P^u_i$ with $P^u_1$ containing the leading term of $P^u_{a_1,\ldots,a_{q_u}}$. For example the element $f_{a_1,\ldots,a_{s_1}}$ in the Lemma 2.5 is written as $f_{a_1,\ldots,a_{s_1}}=\sum_{i=1}^{s_2}f_i$.

Given $P^u_{a_1,\ldots,a_{q_u}}$ and $P^v_{b_1,\ldots,b_{q_v}}$ in $G$, define $m_{12}$, $m_{21}$, $M_{12}$ and $M_{21}$ as definition 4.2. Assume $M_{12}$ has column index $c_1,\ldots,c_{p_{12}}$ and $M_{21}$ has column indices,

$d_1, ..., d_{p_{21}}$. Define $\overline{P^u_{a_1,..,a_{q_u},d_1,...,d_{p_{21}}}}$ and $\overline{P^v_{b_1,...,b_{q_v},c_1,...,c_{p_{12}}}}$ as following: add the rows of $M_{21}$ on top of each matrix of $P^u_{a_1,...,a_{q_u}}$ and add the columns of $M_{21}$ in front of each matrix of $P^u_{a_1,...,a_{q_u}}$. Take the determinant of each matrix and take the sum of all determinants to form the new polynomial $\overline{P^u_{a_1,..,a_{q_u},d_1,...,d_{p_{21}}}}$. Similarly, use $M_{12}$ to obtain $\overline{P^v_{b_1,...,b_{q_v},c_1,...,c_{p_{12}}}}$ . Write $\overline{P^u_{a_1,..,a_{q_u},d_1,...,d_{p_{21}}}} = \sum_{i=1}^{p_u} \overline{P^u_i}$ and $\overline{P^v_{b_1,...,b_{q_v},c_1,...,c_{p_{12}}}} = \sum_{i=1}^{p_v} \overline{P^v_i}$ where $\overline{P^u_1}$ and $\overline{P^v_1}$ contain the leading terms of $\overline{P^u_{a_1,..,a_{q_u},d_1,...,d_{p_{21}}}}$ and $\overline{P^v_{b_1,...,b_{q_v},c_1,...,c_{p_{12}}}}$ . For example in Lemma 2.5, we have $f_{a_1,...,a_{s_1}} = f^1_{a_1,...,a_{s_1}}$ and $f_{b_1,a_2,...,a_{s_1}} = f^2_{b_1a_2,...,a_{s_1}}$ then $m_{12} = M_{12} = z_{a_1}$ and $m_{21} = M_{21} = z_{b_1}$ then

$$\begin{aligned}
\overline{f^1_{b_1,a_1,a_2,...,a_{s_1}}} &= \sum_{q=1}^{s_2} (-1)^{q+1} \left| \begin{bmatrix} Z^{1,1} \\ Z^{q,q} \\ Y^{1,q-1} \\ X^{q+1,m} \end{bmatrix}_{b_1,a_1,a_2,...,a_{s_1}} \right| = \sum_{i=1}^{s_2} \overline{f_i}, \\
&= -\overline{f^2_{a_1,b_1,a_2,...,a_{s_1}}} = -\sum_{i=1}^{s_1} \overline{f^2_i}.
\end{aligned}$$

The technique of proving the following lemma is the main technique we are going to use for computing the $S$-pairs of elements of a Gröbner basis.

**Lemma 4.6.** *Notation as above. If*

$$\operatorname{in}(m_{21}P^u_{a_1,...,a_{q_u}}) = \operatorname{in}(|M_{21}|P^u_{a_1,...,a_{q_u}}) = \operatorname{in}(\overline{P^u_{a_1,..,a_{q_u},d_1,...,d_{p_{21}}}}) = \operatorname{in}(\overline{P^u_1})$$

*and*

$$\operatorname{in}(m_{12}P^v_{b_1,...,b_{q_v}}) = \operatorname{in}(|M_{12}|P^v_{b_1,...,b_{q_v}}) = \operatorname{in}(\overline{P^v_{b_1,...,b_{q_v},c_1,...,c_{p_{12}}}}) = \operatorname{in}(\overline{P^v_1}).$$

*Furthermore* $\sum_{i=2}^{p_u} \overline{P^u_i}$ *and* $\sum_{i=2}^{p_v} \overline{P^v_i}$ *can be written as combination of elements of* $G$ *with leading term smaller than* $\operatorname{in}(\overline{P^u_1})$. *Then the* $S$*-pair of* $P^u_{a_1,...,a_{q_u}}$ *and* $P^v_{b_1,...,b_{q_v}}$ *has zero reminder.*

*Proof.* From the definition of $M_{12}$, $M_{21}$, we have $\overline{P^u_1} = \overline{P^v_1}$. Hence the following equation holds

$$\overline{P^u_{a_1,..,a_{q_u},d_1,...,d_{p_{21}}}} - \overline{P^v_{b_1,...,b_{q_v},c_1,...,c_{p_{12}}}} = \sum_{i=2}^{u} \overline{P^u_i} - \sum_{i=2}^{v} \overline{P^v_i}. \tag{4.1}$$

$\overline{P^u_{a_1,..,a_{q_u},d_1,...,d_{p_{21}}}}$ can be written as

$$\sum_{\{\alpha_1,...,\alpha_{p_{21}}\} \cup \{\beta_1,...,\beta+q_u\} = \{a_1,...,a_{q_u},d_1,...,d_{p_{21}}\}} |M^{21}_{\alpha_1,...,\alpha_{p_{21}}}|P^u_{\beta_1,...,\beta_{q_u}}$$

where $M^{21}_{\alpha_1,...,\alpha_{p_{21}}}$ has the same rows as $M^{21}$ with columns, $\alpha_1, ..., \alpha_{p_{21}}$ and $P^u_{\beta_1,...,\beta_{q_u}}$ is in $G$ with columns, $\beta_1, ..., \beta_{q_u}$. Similarly, $\overline{P^v_{b_1,...,b_{q_v},c_1,...,c_{p_{12}}}}$ can be written as

$$\sum_{\{\alpha_1,...,\alpha_{p_{21}}\}\cup\{\beta_1,...,\beta+q_u\}=\{a_1,...,a_{q_u},d_1,...,d_{p_{21}}\}} |M^{12}_{\alpha_1,...,\alpha_{p_{21}}}|P^v_{\beta_1,...,\beta_{q_u}}.$$

$|M_{21}|P^u_{a_1,...,a_{q_u}}$ and $|M_{12}|P^v_{b_1,...,b_{q_v}}$ are one of summands and their initial terms are the initial terms of each sum. After moving everything other than $m_{21}P^u_{a_1,...,a_{q_u}}$ and $m_{12}P^v_{b_1,...,b_{q_v}}$ from the left-hand side of 4.1 to the right-hand side, we obtain the equality:

$$m_{21}P^u_{a_1,...,a_{q_u}} - m_{12}P^v_{b_1,...,b_{q_v}} = \sum r_i g_i$$

with $g_i \in G$ and $\text{in}(r_i g_i) < \text{in}(m_{12}P^v_{b_1,...,b_{q_v}})$. $\square$

We are going to define some polynomials that are in the ideal $\mathcal{L}$. Those polynomials will be part of the Gröbner basis of $\mathcal{L}$ that we are going to compute. The following definition is coming from the $f_{a_1,...,a_{s_1}}$ as defined in Lemma 2.5.

**Definition 4.7.** Let $1 \le a_1 < a_2 < ... < a_{s_1+k-1} \le \min\{t_1, t_2\}$, and $1 \le l \le k \le s_2$, we define $f^{l,k}_{a_1,...,a_{s_1+k-1}}$ as follow:

$$f^{l,k}_{a_1,...,a_{s_1+k-1}} :=$$

$$\sum_{r=k}^{s_2}(-1)^{r+1}\left|\begin{bmatrix} Z^{l,k-1} \\ Z^{r,r} \\ X^{1,l-1} \\ Y^{1,r-1} \\ Y^{r+1,s_1} \end{bmatrix}_{a_1,...,a_{s_1+k-1}}\right|$$

$$+\sum_{r=k}^{s_2}(-1)^{r+1}\sum_{u=r+1}^{s_1}\left|\begin{bmatrix} Z^{l,k-1} \\ X^{r,r}-Y^{r,r} \\ X^{1,l-1} \\ Y^{1,r-1} \\ Y^{r+1,u-1} \\ Z^{u,u} \\ X^{u+1,s_1} \end{bmatrix}_{a_1,...,a_{s_1+k-1}}\right|.$$

**Lemma 4.8.** $f^{l,k}_{a_1,...,a_{s_1+k-1}} \in \mathcal{L}$.

*Proof.* We first define $p^{l,k}_{a_1,\dots,a_{s_1+k-1}}$ as follows:

$$p^{l,k}_{a_1,\dots,a_{s_1+k-1}} = \sum_{r=k}^{s_2}(-1)^{r+1}\left|\begin{bmatrix} Z^{l,k-1} \\ Z^{r,r} \\ X^{1,l-1} \\ Y^{1,r-1} \\ X^{r+1,s_1} \end{bmatrix}_{a_1,\dots,a_{s_1+k-1}}\right|$$

where $1 \leq a_1 < a_2 < \dots < a_{s_1+k-1} \leq \min\{t_1, t_2\}$, and $1 \leq l \leq k \leq 1$. We notice that $p^{1,1}_{a_1,\dots,a_{s_1}} = f_{a_1,\dots,a_{s_1}}$. We will show $p^{l,k}_{a_1,\dots,a_{s_1+k-1}} \in \mathcal{L}$. Since $p^{l,k+1}_{a_1\dots a_{s_1+k}} = \sum_{i=1}^{s_1+k}(-1)^{i+1}z_{ka_i}f^{l,k}_{a_1\dots\hat{a_i}\dots a_{s_1+k}}$ and $p^{l+1,l+1}_{a_1\dots a_{s_1+l}} = \sum_{i=1}^{s_1+l}(-1)^{i+1}x_{la_i}p^{l,l}_{a_1\dots\hat{a_i}\dots a_{s_1+l}}$, we have that the $p^{l,k}_{a_1\dots a_{s_1+k-1}}$'s are all in $\mathcal{L} \subset \mathcal{J}$. By Lemma 4.3, we have

$$\begin{aligned}
p^{l,k}_{a_1,\dots,a_{s_1}} &= \sum_{r=k}^{s_2}(-1)^{r+1}\left|\begin{bmatrix} Z^{l,k-1} \\ Z^{r,r} \\ X^{1,l-1} \\ Y^{1,r-1} \\ X^{r+1,s_1} \end{bmatrix}_{a_1,\dots,a_{s_1+k-1}}\right| \\
&= \textstyle\sum_{r=k}^{s_2}(-1)^{r+1}\left|\begin{bmatrix} Z^{l,k-1} \\ Z^{r,r} \\ X^{1,l-1} \\ Y^{1,r-1} \\ Y^{r+1,s_1} \end{bmatrix}_{a_1,\dots,a_{s_1+k-1}}\right| \\
&\quad + \textstyle\sum_{r=k}^{s_2}(-1)^{r+1}\sum_{u=r+1}^{s_1}\left|\begin{bmatrix} Z^{l,k-1} \\ X^{r,r}-Y^{r,r} \\ X^{1,l-1} \\ Y^{1,r-1} \\ Y^{r+1,u-1} \\ Z^{u,u} \\ X^{u+1,s_1} \end{bmatrix}_{a_1,\dots,a_{s_1+k-1}}\right| \\
&\quad + \textstyle\sum_{r=k}^{s_2}(-1)^{r+1}\sum_{u=r+1}^{s_1}\sum_{\{c_1,c_2,d_1,\dots,d_{s_1+k-3}\}=\{a_1,\dots,a_{s_1+k-1}\}} \\
&\qquad \left(\pm(g_{rc_1,uc_2}-g_{rc_2,uc_1})\left|\begin{bmatrix} Z^{l,k-1} \\ X^{1,l-1} \\ Y^{1,r-1} \\ Y^{r+1,u-1} \\ X^{u+1,s_1} \end{bmatrix}_{d_1,\dots,d_{s_1+k-3}}\right|\right).
\end{aligned}$$

Since $p^{l,k}_{a_1,\ldots,a_{s_1+k-1}} \in \mathcal{L}$, and

$$\sum_{r=k}^{s_2}(-1)^{r+1}\sum_{u=r+1}^{s_1}\sum_{\{c_1,c_2,d_1,\ldots,d_{s_1+k-3}\}=\{a_1,\ldots,a_{s_1+k-1}\}}$$
$$\pm(g_{rc_1,uc_2}-g_{rc_2,uc_1})\left|\begin{bmatrix} Z^{l,k-1}\\ X^{1,l-1}\\ Y^{1,r-1}\\ Y^{r+1,u-1}\\ X^{u+1,s_1}\end{bmatrix}_{d_1,\ldots,d_{s_1+k-3}}\right| \in \mathcal{L},$$

we have

$$\begin{aligned} f^{l,k}_{a_1,\ldots,a_{s_1+k-1}} &= \sum_{r=k}^{s_2}(-1)^{r+1}\left|\begin{bmatrix} Z^{l,k-1}\\ Z^{r,r}\\ X^{1,l-1}\\ Y^{1,r-1}\\ Y^{r+1,s_1}\end{bmatrix}_{a_1,\ldots,a_{s_1+k-1}}\right| \\ &+\sum_{r=k}^{s_2}(-1)^{r+1}\sum_{u=r+1}^{s_1}\left|\begin{bmatrix} Z^{l,k-1}\\ X^{r,r}-Y^{r,r}\\ X^{1,l-1}\\ Y^{1,r-1}\\ Y^{r+1,u-1}\\ Z^{u,u}\\ X^{u+1,s_1}\end{bmatrix}_{a_1,\ldots,a_{s_1+k-1}}\right| \in \mathcal{L}. \end{aligned}$$

□

The following definition is coming from the $S$-pairs of $X_{a_1,\ldots,a_{s_1}}$ and $g_{ij,lk}$ as defined in the Lemma 2.5.

**Definition 4.9.** Let $1 \leq p_1 \leq m$, $1 \leq q_1 \leq n$, $a_{s_1} < ... < a_j \leq q_1 < a_{j-1} < ... < a_1$. We define $U_{p_1,q_1,a_1,...,a_{s_1}}$ as follows:

$$U_{p_1,q_1,a_{s_1},\ldots,a_1} := z_{p_1q_1}\left|\begin{bmatrix} & & X^{1,p_1-1} & & & \\ x_{p_1a_{s_1}} & \cdots & x_{p_1a_j} & y_{p_1a_{j-1}} & \cdots & y_{p_1a_1} \\ & & Y^{p_1+1,s_1} & & & \end{bmatrix}\right|$$

$$+\sum_{k=j+1}^{m}(x_{p_1q_1}-y_{p_1q_1})(-1)^{k+p_1}z_{p_1a_k}|X^{1,\ldots,\hat{p_1},\ldots,m}_{a_1,\ldots,\hat{a_k},..a_m}|$$

$$+\sum_{u=p_1+1}^{s_1}(x_{p_1q_1}-y_{p_1q_1})\left|\begin{bmatrix} & & X^{1,p_1-1} & & & \\ x_{p_1a_{s_1}} & \cdots & x_{p_1a_j} & y_{p_1a_{j-1}} & \cdots & y_{p_1a_1} \\ & & Y^{p_1+1,u-1} & & & \\ & & Z^{u,u} & & & \\ & & X^{u+1,s_1} & & & \end{bmatrix}_{a_1,\ldots,a_{s_1}}\right|.$$

**Lemma 4.10.** $U_{p_1q_1a_1,\ldots,a_{s_1+k-1}} \in \mathcal{L}$.

*Proof.* We use Lemma 4.3 on $|X^{1,s_1}_{a_1,\ldots,a_{s_1}}|$, then we have

$$\begin{aligned}\alpha = {} & z_{p_1,q_1}|X_{a_1,\ldots,a_{s1}}| \\ = {} & z_{p_1q_1}\left(\left|\begin{bmatrix} & & X^{1,p_1-1} & & & \\ x_{p_1a_{s_1}} & \cdots & x_{p_1a_j} & y_{p_1a_{j-1}} & \cdots & y_{p_1a_1} \\ & & Y^{p_1+1,s_1} & & & \end{bmatrix}\right|\right. \\ & +\sum_{k=j+1}^{m}(-1)^{k+p_1}(x_{p_1a_k}-y_{p_1a_k})|X^{1,\ldots,\hat{p_1},\ldots,m}_{a_1,\ldots,\hat{a_k},\ldots,a_m}| + \sum_{u=p_1+1}^{s_1}\sum_{k=1}^{s_1} \\ & \left.(-1)^{u+k}(x_{ua_k}-y_{ua_k})\left|\begin{bmatrix} & & X^{1,p_1-1} & & & \\ x_{p_1a_{s_1}}\cdots & & x_{p_1a_j} & y_{p_1a_{j-1}}\cdots & & y_{p_1a_1} \\ & & X^{p_1+1,u-1} & & & \\ & & X^{u+1,s_1} & & & \end{bmatrix}_{a_1,\ldots,\widehat{a_k},\ldots,a_m}\right|\right).\end{aligned}$$

Review the definition of $g_{ij,lk}$. We substitute all the monomials that are the leading terms of $\{g_{ij,lk}\}$. The above expression becomes

$$z_{p_1q_1}\left|\begin{bmatrix} X^{1,p_1-1} \\ x_{p_1a_{s_1}}\cdots \quad x_{p_1a_j}\; y_{p_1a_{j-1}}\cdots \quad y_{p_1a_1} \\ Y^{p_1+1,s_1} \end{bmatrix}\right| + \sum_{k=j+1}^{s_1}(-1)^{p_1+k}g_{p_1q_1,p_1a_k}|X^{1,\dots,\hat{p_1},\dots,m}_{a_1,\dots,\hat{a_k},..a_m}|$$

$$+\sum_{k=j+1}^{m}(x_{p_1q_1}-y_{p_1q_1})(-1)^{k+p_1}z_{p_1a_k}|X^{1,\dots,\hat{p_1},\dots,m}_{a_1,\dots,\hat{a_k},..a_m}|$$

$$+\sum_{u=p_1+1}^{s_1}\sum_{k=1}^{m}(-1)^{k+u}(g_{p_1q_1,u,a_k}-g_{p_1a_k,u,q_1})$$

$$\left(\left|\begin{bmatrix} X^{1,p_1-1} \\ x_{p_1a_{s_1}}\cdots \quad x_{p_1a_j}\; y_{p_1a_{j-1}}\cdots \quad y_{p_1a_1} \\ Y^{p_1+1,u-1} \\ X^{u+1,s_1} \end{bmatrix}_{a_1,\dots,\widehat{a_k},\dots,a_m}\right|\right)$$

$$+\sum_{u=p_1+1}^{s_1}(x_{p_1q_1}-y_{p_1q_1})\left|\begin{bmatrix} X^{1,p_1-1} \\ x_{p_1a_{s_1}} \quad \cdots \quad x_{p_1a_j} \quad y_{p_1a_{j-1}} \quad \cdots \quad y_{p_1a_1} \\ Y^{p_1+1,u-1} \\ Z^{u,u} \\ X^{u+1,s_1} \end{bmatrix}_{a_1,\dots,a_{s_1}}\right|.$$

We define $\beta$ as follows:

$$\beta = \sum_{k=j+1}^{s_1}(-1)^{p_1+k}g_{p_1q_1,p_1a_k}|X^{1,\dots,\hat{p_1},\dots,m}_{a_1,\dots,\hat{a_k},..a_m}| + \sum_{u=p_1+1}^{s_1}\sum_{k=1}^{m}(-1)^{k+u}(g_{p_1q_1,u,a_k}-g_{p_1a_k,u,q_1})$$

$$\left(\left|\begin{bmatrix} X^{1,p_1-1} \\ x_{p_1a_{s_1}}\cdots \quad x_{p_1a_j}\; y_{p_1a_{j-1}}\cdots \quad y_{p_1a_1} \\ Y^{p_1+1,u-1} \\ X^{u+1,s_1} \end{bmatrix}_{a_1,\dots,\widehat{a_k},\dots,a_m}\right|\right).$$

$\beta$ is in $\mathcal{L}$ and $\alpha$ is in $\mathcal{L}$, hence $U_{p_1q_1a_1,\dots,a_{s_1}} = \alpha - \beta$ is in $\mathcal{L}$. □

The following definition is coming from the $S$-pairs of $U_{p,q,a_1,\dots,a_{s_1}}$ as defined in the Definition 4.9 and $Y_{a_1,\dots,a_{s_2}}$ as defined in the Lemma 2.5.

**Definition 4.11.** Let $1 \le b_{s_2} < \dots < b_1 \le n$, $1 \le p_1 \le m$, $1 \le q_1 \le n$, $a_{s_1} < \dots < a_{s_2+1} < a_{p_1} < \dots < a_1$ and $a_{p_1} \le q_1$. Let $i$ be integer so that $1 \le i \le p$ and $a_{s_2+1} < b_{s_2} < \dots < b_{i+1} < a_{p_1-1} \le b_i$ and $b_l \ne a_{p_1}$ for $l \ge i+1$.

We define $M_{12}$ as follows:

$$M_{12} = z_{p_1q_1}x_{p_1a_{p_1}}\left|\begin{bmatrix} X^{1,p_1-1} \\ Y^{s_2+1,s_1} \end{bmatrix}_{a_1,\dots,a_{p_{1}-1},a_{s_2+1},\dots,a_{s_1}}\right|.$$

We define

$$\begin{gathered} W_{p_1,q_1,a_1,\dots,a_{p1},a_{s_2+1},\dots,a_{s_1},b_1,\dots,b_{s_2}} := \\ M_{12}|Y^{1,s_1}_{b_1,\dots,b_{s_2}}| - \left|Y^{1,i}_{b_1,..,b_i}\right| \sum_{\{c_{i+1},\dots,c_{p_1},d_{p_1+1},\dots,d_{s_2}\}=\{b_{i+1},\dots.,b_{s_2}\}} \\ \left|Y^{i+1,p_1}_{c_{i+1},..,c_{p_1}}\right| U_{p_1q_1,a_1,\dots,a_{p_1-1},a_{p_1},d_{p_1+1},\dots,d_{s_2},a_{s_2+1},\dots,a_{s_1}}. \end{gathered}$$

*Remark* 4.12. From the way we define $W_{p_1,q_1,a_1,\dots,a_{p_1},a_{s_2+1},\dots,a_{s_1},b_1,\dots,b_{s_1}}$, it is in $\mathcal{L}$. Notice that all the submatrices $|Y^{p_1,s_2}_{d_{p_1+1},\dots,d_{s_2}}|$ of $|Y^{1,s_2}_{b_1,\dots,b_{s_2}}|$ such that $a_{s_2+1} < d_{s_2} < \dots < d_{p_1+1} < a_{p_1-1}$ are cancelled. Hence the leading term is

$$\mathrm{in}(M_{12}|Y^{1,i-1}_{b_1,\dots,b_{i-1}}||Y^{i,p_1}_{b_{i+1},\dots,b_{p_1+1}}||Y^{p_1+1,s_2}_{b_{s_2},\dots,b_{p_1+2},b_i}|).$$

The following definition is coming from the $S$-pairs of $W_{p,q,a_1,\dots,a_p,a_{s_2+1},\dots,a_{s_1},b_1,\dots,b_{s_1}}$ as defined in the Definition 4.11 and $U_{p,q,a_1,\dots,a_{s_1}}$ as defined in the Definition 4.9.

**Definition 4.13.** Let $1 \le p_1 \le m$, $1 \le q_1 \le n$, $v = p_1+1, \dots, s_2-1$, $1 \le a_{s_1} < \dots < a_{s_2+1} < a_{p_1} < \dots < a_1 \le t_1$ and $a_{p_1} \le q_1$. Let $i$ be integer so that $1 \le i \le p$ and let $a_{s_2+1} < b_{s_2} < \dots < b_{v+2} < b'_v < \dots < b'_{p_1+1} < b_{p_1} < \dots < b_{i+1} < a_{p_1-1} \le b_{p_1+1}$ and $b'_l \ne a_{p_1}$ for $l \ge i+1$ and $b'_{v-1} \le b_{v+1}$. Let $a_{s_1} < \dots. < a_{s_2+1} < b_{s_2} < \dots < b_{v+2} < b_{v+1} < b_v < b_{v-1} < \dots < b_{p_1+2} < a_{p_1} < a_{p_1-1} \le b_{p_1+1}$, and $b'_r \le b_{r+2} < b_{r+1}$ for $r = p_1, \dots, v-2$.

We define

$$\begin{gathered} W^{p_1+1,v}_{p_1,q_1,a_1,\dots,a_{p1},a_{s_2+1},\dots,a_{s_1},b_1,\dots,b_{p_1+1},b_{p_1+2},b_{p_1+3}.,b_{s_2},b'_{p_1+1},b'_{p_1+2},\dots,b'_v} := \\ y_{v-1,b'_{v-1}}y_{v,b_v}W^{p_1+1,v-2}_{p_1,q_1,a_1,\dots,a_{p1},a_{s_2+1},\dots,a_{s_1},b_1,\dots,b_{v-1},b'_v,b_{v+1},\dots,b_{s_2},b'_{p_1+1},b'_{p_1+2},\dots,b'_{v-2}} \\ -y_{v,b'_v}W^{p_1+1,v-1}_{p_1,q_1,a_1,\dots,a_{p1},a_{s_2+1},\dots,a_{s_1},b_1,\dots,b_{s_2},b'_{p_1+1},b'_{p_1+2},\dots,b'_{v-1}}. \end{gathered}$$

Here

$$W^{p_1+1,p_1-1}_{p_1,q_1,a_1,\dots,a_{p_1},a_{s_2+1},\dots,a_{s_1},b_1,\dots,b_{s_2}} = U_{p_1,q_1,a_1,\dots,a_{s_1}}$$

and

$$W^{p_1+1,p_1}_{p_1q_1,a_1,\dots,a_{p_1},a_{s_2+1},\dots,a_{s_1},b_1,\dots,b_{s_2}} = W_{p_1q_1,a_1,..,a_{p_1},a_{s_2+1},\dots,a_{s_1},b_1,\dots,b_{s_2}}.$$

*Remark* 4.14. From the way we define $W^{p_1+1,v}_{p_1,q_1,a_1,\dots,a_{p_1},a_{s_2+1},\dots,a_{s_1},b_1,\dots,b_{s_2},b'_{p_1+1},\dots,b'_v}$, it is clear that it belongs to $\mathcal{L}$. Notice it has leading term

$$\operatorname{in}(z_{p_1q_1}\left|\begin{bmatrix} X^{1,p_1} \\ Y^{s_2+1,s_1} \end{bmatrix}_{a_1,\dots,a_{p_1},a_{s_2+1},\dots,a_{s_1}}\right| |Y^{1,i-1}_{b_1,\dots,b_{i-1}}||Y^{i,p_1}_{b_{i+1},\dots,b_{p_1+1}}|$$
$$y_{p_1+1,b_i}y_{p_1+1,b_{p_1+1}}y_{p_1+2,b_{p_1+2}}y_{p_1+2,b'_{p_1+2}}\cdots y_{v-1,b'_{v-1}}y_{v-1,b'_{v-1}}y_{vb_v}y_{v,b'_v}$$
$$|Y^{l+1,s_2}_{b_{s_2},\dots,b_{v+2},b_{v+1}}|).$$

The following definition is coming from the $S$-pairs of $f^{l,k}_{a_1,\dots,a_{s_1+k-1}}$ as defined in the Definition 4.7 and $Y_{a_1,\dots,a_{s_2}}$ as defined in the Definition 2.5.

**Definition 4.15.** Let $b_{s_1} < \dots < b_1$, and $1 \le p_l < \dots < p_k < b_{s_1} < \dots < b_{s_2+1} < c_{s_2} < \dots < c_{k+1} < b_{k-1} < \dots < b_1 < a_{k-1} < \dots < a_1 \le t_1$.

Let

$$M_{12} = \left|\begin{bmatrix} Z^{l,k} \\ X^{1,k-1} \\ Y^{s_2+1,s_1} \end{bmatrix}_{p_l,..,p_k,a_1,\dots,a_{k-1},b_{s_2+1},\dots,b_{s_1}}\right|.$$

We define

$$V_{p_l,\dots,p_k,a_1,\dots,a_{k-1},b_1,\dots,b_{s_1}} :=$$
$$M_{12}|Y^{1,s_2}_{b_1,\dots,b_{s_2}}| - \sum_{\{e_k,c_{k+1},\dots,c_{s_2}\}=\{b_k,\dots.,b_{s_2}\}}$$
$$\pm y_{ke_k} f^{l,k}_{p_l,\dots,p_k,a_1,\dots,a_{k-1},b_1,\dots,b_{k-1},c_{k+1},\dots,c_{s_2},b_{s_2+1},\dots,b_{s_1}}.$$

*Remark* 4.16. From the way we define $V_{p_l,\dots,p_k,a_1,\dots,a_{k-1},b_1,\dots,b_{s_1}}$, it is in $\mathcal{L}$. Notice the submatrices $|Y^{k+1,s_2}_{c_{k+1},\dots,c_{s_2}}|$ of $|Y^{1,s_2}_{b_1,\dots,b_{s_2}}|$ such that $b_{s_2+1} < c_{s_2} < \dots < c_{k+1} < b_{k-1}$ are cancelled. Hence the leading term of $V_{p_l,\dots,p_k,a_1,\dots,a_{k-1},b_1,\dots,b_{s_1}}$ is

$$\operatorname{in}(M_{12}|Y^{1,k-2}_{b_1,..,b_{k-2}}|y_{k-1,b_k}|Y^{k,s_2}_{b_{s_2},\dots,b_{k+1},b_{k-1}}|).$$

The following definition is coming from the $S$-pairs of elements in $\{V_{p_1,..,p_k,a_1,\dots,a_{k-1},b_1,\dots,b_{s_1}}\}$ as defined in the Definition 4.15.

**Definition 4.17.** Let $1 \le l \le k \le s_2$ and $1 \le p_l < \dots < p_k < b_{s_1} < \dots < b_{s_2+1} < \dots < b_{k+1} < b_{k-1} < b_k < b_{k-2} < \dots < b_1 < a_{l-1} < \dots < a_1 \le t_1$. Let $w = k, \dots, s_2 - 1$ and $1 \le b_{s_2} < \dots < b_{w+2} < b'_w < b'_{w-1} < \dots < b'_k < b_{k-1} < b_{k-2} \dots < b_1 \le t_2$ and $b'_{w-1} \le b_{w+1}$, and $b'_r \le b_{r+2} < b_{r+1}$ for $r = k, \dots, l-2$.

We define

$$\begin{aligned} &V^{k,w}_{p_l,\dots,p_k,a_1,\dots,a_{k-1},b_1,\dots,b_{s_1},b'_k,b'_{k+1},\dots,b'_w} := \\ &y_{w-1,b_{w-1}}y_{w,b_w}V^{k,w-2}_{p_l,\dots,p_k,a_1,\dots,a_{k-1},b_1,..,b'_w,\dots,b_{s_1},b'_k,b'_{k+1},\dots,b'_{w-2}} \\ &-y_{wb'_w}V^{k,w-1}_{p_l,\dots,p_k,a_1,\dots,a_{k-1},b_1,\dots,b_{s_1},b'_k,b'_{k+1},\dots,b'_{w-1}} . \end{aligned}$$

Here $V^{k,k-2}_{p_l,\dots,p_k,a_1,\dots,a_{k-1},b_1,\dots,b_{s_1}} = V^{k,k-1}_{p_l,\dots,p_k,a_1,\dots,a_{k-1},b_1,\dots,b_{s_1}} = V_{p_l,\dots,p_k,a_1,\dots,a_{k-1},b_1,\dots,b_{s_1}}$.

*Remark* 4.18. From the way we define $V^{k,w}_{p_l,\dots,p_k,a_1,\dots,a_{k-1},b_1,\dots,b_{s_1},b'_k,b'_{k+1},\dots,b'_l}$, it is in $\mathcal{L}$. It has leading term

$$\mathrm{in}(M_{12}|Y^{1,k-2}_{b_1,..,b_{k-2}}|y_{k-1,b_{k-1}}y_{kb_k}y_{kb'_k}...y_{lb_w}y_{lb'_w}|Y^{l+1,s_2}_{b_{s_2},\dots,b_{w+1}}|).$$

The following definition is coming from the $S$-pairs of $g_{ij,lk}$ as defined in the Definition 2.5 and $f^{l,k}_{a_1,\dots,a_{s_1+k-1}}$ as defined in the Definition 4.7.

**Definition 4.19.** Let $1 \le l \le k \le s_2$, $1 \le q \le n$, $1 \le a_{s_1+k-1} < .... < a_1 \le t_1$, $a_{s_1+k-1} < q$, $a_{j+1} \le q < a_j$ for some $j = l-1, ..., s_1+k-3$. Let $\overline{f^{l,k,x_{l-1}}_{a_1,\dots,\hat{a_c},\dots,a_{s_1+k-1}}}$ be the determinant of matrices that coming from deleting row $x_{l-1}$ and column $a_c$. We define $H^{l,k,q}_{a_1,\dots,a_{s_1+k-1}}$ as follows

$$H^{l,k,q}_{a_1,\dots,a_{s_1+k-1}} \quad = \quad z_{l-1,q}f^{l,k}_{a_1,\dots,a_{s_1+k-1}} - \sum_{c=k}^{j}(-1)^{k+c}g_{l-1,q,l-1,a_c}\overline{f^{l,k,x_{l-1}}_{a_1,\dots,\hat{a_c},\dots,a_{s_1+k-1}}}.$$

*Remark* 4.20. It is clear that $H^{l,k,q}_{a_1,\dots,a_{s_1+k-1}}$ is in $\mathcal{L}$ from the way we define it. Notice in the row $x_l$ of $f^{l,k}_{a_1,\dots,a_{s_1+k-1}}$, the $x_{l,a_c}$ are cancelled by the $g_{l-1,q,l-1a_c}$. Hence the leading term of $H^{l,k,q}_{a_1,\dots,a_{s_1+k-1}}$ is

$$z_{l-1,q}\mathrm{in}\left(\left|Z^{l,k}_{p_l,\dots,p_k}\right| x_{l-1,a_{j+1}} \left|X^{1,l-2}_{a_1,\dots,a_{l-2}}\right| \left|\begin{bmatrix} Y^{1,k-1} \\ Y^{k+1,s_1} \end{bmatrix}_{b_1,\dots,b_{k-1},b_{k+1},\dots,b_{s_1}}\right|\right).$$

Here $p_i \neq a_{j+1}$, $b_i \neq a_{j+1}$ for all $i$.

The following definition is coming from the $S$-pairs of $H^{l,k,q}_{a_1,\dots,a_{s_1+k-1}}$ as defined in the Definition 4.19 and $Y_{a_1,\dots,a_{s_2}}$ as defined in the Definition 2.5.

**Definition 4.21.** Let $1 \le l \le k \le s_2$, $1 \le q \le n$, $1 \le a_{s_1+k-1} < .... < a_1 \le t_1$, $a_{s_1+k-1} < q$, $a_{j+1} \le q < a_j$ for some $j = l-1, ..., s_1+k-3$. Let $a_{l+s_2-1} < b_{s_2} < ... < b_k < a_{l-1+k-1} = b_{k-1} < .... < a_{l-1+1} = b_1$.

Let

$$M = z_{l-1,q}x_{l-1,a_{j+1}}\left|\begin{bmatrix} Z^{l,k} \\ X^{1,l-2} \\ Y^{s_2+1,s_1} \end{bmatrix}_{a_{s_1+k-1},\dots,a_{l+s_2-1},a_{l-1},\dots,a_1}\right|.$$

We define

$$\begin{aligned}
&I^{l,k,q}_{a_{s_1+k-1},\dots,a_{l+s_2-1},a_{l-1},\dots,a_1,b_1,\dots,b_{s_2}} := \\
&M|Y^{1,s_2}_{b_1,\dots,b_{s_2}}| - \sum_{\{e_k,c_{k+1},\dots,c_{s_2}\}=\{b_k,\dots,b_{s_2}\}} \\
&\pm y_{ke_k} H^{l,k,q}_{a_{s_1+k-1},\dots,a_{l+s_2-1},c_{s_2},\dots,c_{k+1},b_{k-1},\dots,b_1,a_{l-1},\dots,a_1}.
\end{aligned}$$

*Remark* 4.22. It is clear that $I^{l,k,q}_{a_{s_1+k-1},\dots,a_{l+s_2-1},a_{l-1},\dots,a_1,b_1,\dots,b_{s_2}}$ is in $\mathcal{L}$ from the way we define it. Notice that the submatrices $|Y^{k+1,s_2}_{c_{k+1},\dots,c_{s_2}}|$ of $|Y^{1,s_2}_{b_1,\dots,b_{s_2}}|$ with $a_{l+s_2-1} < c_{s_2} < \dots < c_{k+1} < b_{k-1}$ are cancelled by $H^{l,k,q}$'s, hence the leading term of
$I^{l,k,q}_{a_{s_1+k-1},\dots,a_{l+s_2-1},a_{l-1},\dots,a_1,b_1,\dots,b_{s_2}}$ is

$$\text{in}\left( z_{l-1,q} x_{l-1,a_{j+1}} \left| \begin{bmatrix} Z^{l,k} \\ X^{1,l-2} \\ Y^{s_2+1,s_1} \end{bmatrix}_{a_{s_1+k-1},\dots,a_{l+s_2-1},a_{l-1},\dots,a_1} \right| \right.$$
$$\left. y_{k-1,b_k} \left| \begin{bmatrix} Y^{1,k-2} \\ Y^{k,s_2} \end{bmatrix}_{b_{s_2},\dots,b_{k+1},b_{k-1},b_{k-2}\dots b_1} \right| \right).$$

The following definition is coming from the $S$-pairs of elements of $\{I^{l,k,q}_{q_l,\dots,q_k,b_{s_1},\dots,b_{s_2},\dots,b_1,a_{l-1},\dots,a_1}\}$.

**Definition 4.23.** Let $1 \le l \le k \le s_2$, $1 \le q \le n$, $k \le w \le s_2 - 1$, $1 \le q_l < \dots < q_k < b_{s_1} < \dots < b_{s_2} < \dots < b_{k+1} < b_{k-1} < b_k < b_{k-2} < \dots < b_1 < a_{l-2} < \dots < a_1 \le t_1$, $q_l < q$, $q_l < a_{l-1} \le q$ . Let $w = k, \dots, s_2 - 1$ and $1 \le b_{s_2} < \dots < b_{w+2} < b'_w < b'_{w-1} < \dots < b'_k < b_{k-1} < b_{k-2} \dots < b_1 \le t_2$ and $b'_{w-1} \le b_{w+1}$, and $b'_r \le b_{r+2} < b_{r+1}$ for $r = k, \dots, l-2$.

We define

$$\begin{aligned}
&{}^{k,w}I^{l,k,q}_{q_l,\dots,q_k,b_{s_1},\dots,b_{s_2},\dots,b_1,a_{l-1},\dots,a_1,b'_k,\dots,b'_w} := \\
&y_{w-1,b'_{w-1}} y_{wb_w} {}^{k,w-2}I^{l,k,q}_{q_l,\dots,q_k,b_{s_1},\dots,b_{s_2},\dots,b'_w,\dots,b_1,a_{l-1},\dots,a_1,b'_k,\dots,b'_{w-2}} \\
&-y_{wb'_w} {}^{k,w-1}I^{l,k,q}_{q_l,\dots,q_k,b_{s_1},\dots,b_{s_2},\dots,b_1,a_{l-1},\dots,a_1,b'_k,\dots,b'_{w-1}}.
\end{aligned}$$

Here

$$\begin{aligned}
{}^{k,k-2}I^{l,k,q}_{q_l,\dots,q_k,b_{s_1},\dots,b_{s_2},\dots,b_1,a_{l-1},\dots,a_1} &= {}^{k,k-1}I^{l,k,q}_{q_l,\dots,q_k,b_{s_1},\dots,b_{s_2},\dots,b_1,a_{l-1},\dots,a_1} \\
= I^{l,k,q}_{q_l,\dots,q_k,b_{s_1},\dots,b_{s_2},\dots,b_1,a_{l-1},\dots,a_1} & .
\end{aligned}$$

*Remark* 4.24. From the way we define ${}^{k,w}H^{l,k,q}_{a_{s_1+k1-},\dots,a_{l-1+s_2}b_{s_2},\dots,b_1,a_{l-1},\dots,a_1,b'_k,\dots,b'_w}$, it is in $\mathcal{L}$. The leading term of ${}^{k,w}H^{l,k,q}_{a_{s_1+k1-},\dots,a_{l-1+s_2}b_{s_2},\dots,b_1,a_{l-1},\dots,a_1,b'_k,\dots,b'_w}$ is

$$\text{in}\left( z_{l-1,q}x_{l-1,a_{j+1}} \left| \begin{bmatrix} Z^{l,k} \\ X^{1,l-2} \\ Y^{s_2+1,s_1} \end{bmatrix}_{a_{s_1+k-1},\dots,a_{l+s_2-1},a_{l-1},\dots,a_1} \right| \right.$$
$$\left. y_{k-1,b_{k-1}}y_{kb_k}y_{kb'_k}\cdots y_{wb_w}y_{wb'_w} \left| \begin{bmatrix} Y^{1,k-2} \\ Y^{w+1,s_2} \end{bmatrix}_{b_{s_2},\dots,,b_{w+1},b_{k-2}\dots b_1} \right| \right).$$

We are now ready to find the Gröbner basis of $\mathcal{L}$.

**Theorem 4.25.** *Use the notation of Definition 4.7, 4.9, 4.11, 4.13, 4.15, 4.17, 4.19, 4.21, 4.23 and let* $\mathcal{G} := \{|X^{1,s_1}_{a_1,..,a_{s_1}}|,\ |Y^{1,s_2}_{b_1,\dots,b_{s_2}}|,\ g_{p_1q_1,p_2q_2},\ f^{l,k}_{a_1,\dots,a_{s_1+k-1}},$
$U_{p_1,q_1,a_1,\dots,a_{s_1}}$ ,
$W_{p,q,a_1,\dots,a_p,a_{s_2+1},\dots,a_{s_1},b_1,\dots,b_{s_2}}$,
$W^{p_1+1,v}_{p_1,q_1,a_1,\dots,a_{p1},a_{s_2+1},\dots,a_{s_1},b_1,\dots,b_{p_1+1},b_{p_1+2},b_{p_1+3}.,b_{s_2},b'_{p_1+1},b'_{p_1+2},\dots,b'_v}$,
$V_{p_l,\dots,p_k,a_1,\dots,a_{k-1},b_1,\dots,b_{s_1}}$, $V^{k,w}_{p_l,\dots,p_k,a_1,\dots,a_{k-1},b_1,\dots,b_{s_1},b'_k,b'_{k+1},\dots,b'_w}$, $H^{l,k,q}_{a_1,\dots,a_{s_1+k-1}}$ ,
$I^{l,k,q}_{a_{s_1+k-1},\dots,a_{l+s_2-1},a_{l-1},\dots,a_1,b_1,\dots,b_{s_2}}$, ${}^{k,w}I^{l,k,q}_{q_l,\dots,q_k,b_{s_1},\dots,b_{s_2},\dots,b_1,a_{l-1},\dots,a_1,b'_k,\dots,b'_w}\}$.

*The* $\mathcal{G}$ *is a Gröbner basis of* $\mathcal{L}$ *with respect to the lexicographic term order and the variables ordered by* $z_{ij} > x_{lk} > y_{pq}$ *for any* $i, j, l, k, p, q$ *and* $x_{ij} < x_{lk}$, $y_{ij} < y_{lk}$ *if* $i > l$ *or* $i = l$ *and* $j < k$ *and* $z_{ij} < z_{lk}$ *if* $i > l$ *or if* $i = l$ *and* $j > k$.

We break up the proof of the above theorem into a sequence of lemmas when we treat $S$-pairs between elements of $\mathcal{G}$. We only have to compute the $S$-pairs of elements whose leading terms are not relative prime. In each lemma, we show $h_{P,Q} = 0$ for some $P, Q$ in $\mathcal{G}$. We define a order on pair $(i, j)$ with $1 \le i \le m$, $1 \le j \le n$. We say $(i, j) > (l, k)$ if $i < l$ or $i = l$ and $j < k$. This is a total order.

**Lemma 4.26.** $h_{P,Q} = 0$ *when* $P$ *and* $Q$ *are in the same group of* $\mathcal{G}$.

*Proof.* We use notation in the Definition 4.5. Notice that $m_{12}$ and $m_{21}$ have the same row indices, $P^u_{a_1,\dots,a_{q_u}}$ and $Q^v_{b_1,\dots,b_{q_v}}$ have the same number of columns, i.e. $q_u = q_v$ hence

$$\overline{P^u_{a_1,..,a_{q_u},d_1,\dots,d_{p_{21}}}} = \overline{Q^v_{b_1,\dots,b_{q_v},c_1,\dots,c_{p_{12}}}}.$$

Also $\text{in}(m_{12}P^u_{a_1,\dots,a_{q_u}}) = \text{in}(m_{21}Q^v_{b_1,\dots,b_{q_u}})$ are indeed the leading terms of $\overline{P^u_{a_1,..,a_{q_u},d_1,\dots,d_{p_{21}}}}$. The first matrix of $\overline{P^u_{a_1,..,a_{q_u},d_1,\dots,d_{p_{21}}}}$ has determinant equal to zero, since it has repeated row. Hence we have $m_{12}P^u_{a_1,\dots,a_{q_u}}$ and $m_{21}P^v_{b_1,\dots,b_{q_u}}$ having different signs in the sum. Except $P^u_{a_1,\dots,a_{q_u}} = f^{l,k}_{a_1,\dots,a_{q_u}}$ and $Q^v_{b_1,\dots,b_{q_v}} = f^{l,k}_{a_1,\dots,b_{q_u}}$ with $a_i = b_i$ for $i \ne k-l+1$ and $a_{k-l+1} \ne b_{k-l+1}$, each summand of all possible cases will have either repeated row, or all the rows, $y_1, ..., y_{s_2}$ or rows of lemma 4.4. Hence they give

$$\sum_{i=2}^{u} \overline{P_i} \in \mathcal{G}.$$

For the remaining case,

$$\sum_{i=2}^{u} \overline{P_i} = f^{l,k+1}_{a_1,...,a_{k-l+1},b_{k-l+1},a_{k-l+2},...,a_{q_u}}$$

from the proof of Lemma 4.8. Hence the following is true:

$$\overline{P^u_{a_1,..,a_{q_u},d_1,...,d_{p_{21}}}} = \sum_{i=2}^{u} \overline{P_i} \in \mathcal{G}.$$

After moving everything other than $m_{12}P^u_{a_1,...,a_{q_u}}$ and $m_{21}Q^v_{b_1,...,b_{q_v}}$ from the left hand side to the right hand side, we obtain the $S$-pair and it becomes:

$$m_{12}P^u_{a_1,...,a_{q_u}} - m_{21}Q^v_{b_1,...,b_{q_v}} = \sum r_i f_i$$

with $\text{in}(r_i f_i) < \text{in}(m_{12}P^u_{a_1,...,a_{q_u}})$ and $f_i \in \mathcal{G}$. □

**Lemma 4.27.** *$h_{P,Q} = 0$ when $P \in \{|X^{1,s_1}_{a_1,...,a_{s_1}}|\}$ in $\mathcal{G}$.*

*Proof.* As the notation in Definition 4.5, we look at $\sum_{i=2}^{u} \overline{P_i}$. For most of cases, $\sum_{i=2}^{u} \overline{P_i} = 0$, since each summand has repeated rows $x_j$ for some $j = 1, ..., s_1$. In some other cases, we have either rows, $y_i$, $x_i$ and $z_i$ in each matrix then Lemma 3.2 can be applied. Or the part of the sum has sum as Lemma 4.4 then deduce that it is in $(\{|Y^{1,s_2}_{b_1,...,b_{s_2}}|\}) + (\{g_{ij,lk}\})$. Similarly, $\sum_{i=2}^{u} \overline{Q_i} \in (\{|Y^{1,s_2}_{b_1,...,b_{s_2}}|\}) + (\{g_{ij,lk}\})$, hence Lemma 4.6 applies. □

**Lemma 4.28.** *$h_{P,Q} = 0$ when $P \in \{|Y^{1,s_2}_{b_1,...,b_{s_2}}|\}$ in $\mathcal{G}$.*

*Proof.* The computation of $S$-pair between $f^{l,k}_{a_1,...,a_{s_1+k-1}}$ and $|Y^{1,s_2}_{b_1,...,b_{s_2}}|$ gives us $V_{p_l,...,p_k,a_1,...,a_{k-1},b_1,...,b_{s_1}}$ as in Definition 4.15. The $S$-pair between
$V_{p_l,...,p_k,a_1,...,a_{k-1},b_1,...,b_{s_1}}$ and $|Y^{1,s_1}_{b_1,...,b_{s_2}}|$ gives us $V^{k,w}_{p_l,...,p_k,a_1,...,a_{k-1},b_1,...,b_{s_1},b'_k,b'_{k+1},...,b'_w}$ as Definition 4.17. The $S$-pair between $V^{k,w}_{p_l,...,p_k,a_1,...,a_{k-1},b_1,...,b_{s_1},b'_k,b'_{k+1},...,b'_w}$ and $|Y^{1,s_2}_{b_1,...b_{s_2}}|$ gives $V^{k,w+1}_{p_l,...,p_k,a_1,...,a_{k-1},b_1,...,b_{s_1},b'_k,b'_{k+1},...,b'_{w+1}}$. Similarly the computation of $S$-pair between $H^{l,k,q}_{a_1,...,a_{s_1+k-1}}$ and $|Y^{1,s_2}_{b_1,...,b_{s_2}}|$ gives $I^{l,k,q}_{a_{s_1+k-1},...,a_{l+s_2-1},a_{l-1},...,a_1,b_1,...,b_{s_2}}$ as Definition 4.21 and the $S$-pair between $I^{l,k,q}_{a_{s_1+k-1},...,a_{l+s_2-1},a_{l-1},...,a_1,b_1,...,b_{s_2}}$ and $|Y^{1,s_2}_{b_1,...,b_{s_2}}|$ gives ${}^{k,w}I^{l,k,q}_{q_l,...,q_k,b_{s_1},...,b_{s_2},...,b_1,a_{l-1},...,a_1,b'_k,...,b'_w}$. Also the $S$-pair between $U_{p_1,q_1,a_1,...,a_{s_1}}$ and $|Y^{1,s_2}_{b_1,...,b_{s_2}}|$ gives $W_{p,q,a_1,...,a_p,a_{s_2+1},...,a_{s_1},b_1,...,b_{s_2}}$ as Definition 4.11 and the $S$-pair between $W_{p,q,a_1,...,a_p,a_{s_2+1},...,a_{s_1},b_1,...,b_{s_2}}$ and $|Y^{1,s_2}_{b_1,...,b_{s_2}}|$ gives
$W^{p_1+1,v}_{p_1,q_1,a_1,...,a_{p1},a_{s_2+1},...,a_{s_1},b_1,...,b_{p_1+1},b_{p_1+2},b_{p_1+3}.,b_{s_2},b'_{p_1+1},b'_{p_1+2},...,b'_v}$. □

**Lemma 4.29.** *$h_{P,Q} = 0$ when $P \in \{g_{ij,lk}\}$ in $\mathcal{G}$.*

*Proof.* If $Q \in \{g_{ij,lk}\}$, we have $Q = z_{p_1q_1}(x_{p_2q_2} - y_{p_2q_2}) - z_{p_2q_2}(x_{p_1q_1} - y_{p_1q_1})$ and $P = g_{ij,lk} = z_{ij}(x_{lk} - y_{lk}) - z_{lk}(x_{ij} - y_{ij})$. It's sufficient to consider either $(p_1, q_1) = (i, j)$ or $(p_2, q_2) = (l, k)$. For the first case,

$$(x_{lk} - y_{lk})Q - (x_{p_2q_2} - y_{p_2q_2})P \quad = (x_{ij} - y_{ij}) \quad g_{lk,p_2q_2}.$$

For the second case,

$$z_{ij}Q - z_{p_1q_1}P = z_{p_2q_2}g_{p_1q_1,ij}.$$

Notice that $P = g_{ij,lk} = z_{i,j}(x_{l,k} - y_{l,k}) - z_{l,k}(x_{i,j} - y_{i,j})$ with $(i, j) > (l, k)$. If $Q \in \{|X^{1,s_1}_{a_1,\ldots,a_{s_1}}|\}$, the computing of $S$-pair of $P$ and $Q$ is similar to Lemma 4.10. And it gives $U_{pqa_1,\ldots,a_{s_1}}$ as Definition 4.9. If $Q \in \{f^{l,k}_{a_1,\ldots,a_{s_{1+k-1}}}\}$, the computing of $S$-pair of $P$ and $Q$ will give us $H^{l,k,q}_{a_1,\ldots,a_{s_1+k-1}}$ as Definition 4.19 when $P = g_{ij,lk}$ and $i = l - 1$. Otherwise $GCD(\text{in}(P),\ \text{in}(Q)) = z_{i,j}$ with $i \in \{l, l + 1, \ldots, k\}$ or $GCD(\text{in}(P),\ \text{in}(Q)) = x_{l,k}$ with $i < l - 1$. For $GCD(\text{in}(P),\ \text{in}(Q)) = z_{i,j}$ with $i \in \{l, l+1, \ldots, k\}$, the computation of $S$-pair gives us $f^{l+1,k}$. For $GCD(\text{in}(P), \text{in}(Q)) = x_{l,k}$, the computation of $S$-pair gives us repeated row, $y_l$, in every matrix of $Q$ and this makes the determinant equal to zero. For all other cases, $Q \in \mathcal{G}$, they come from the $S$-pair of $P \in \{g_{ij,lk}\}$ and $|X^{1,s_1}_{a_1,\ldots,a_{s_1}}|$ or $f^{l,k}_{a_1,\ldots,a_{s_1+k-1}}$. Hence the computations of $S$-pair are very similar. □

**Lemma 4.30.** *$h_{P,Q} = 0$ when $P = f^{l_1,k_1}_{a_1,\ldots,a_{s_1+k_1-1}}$ and $Q = f^{l_2,k_2}_{b_1,\ldots,b_{s_1+k_2-1}}$ and $l_1 \neq l_2$ or $k_1 \neq k_2$ in $\mathcal{G}$.*

*Proof.* We prove this part in two cases: (a) $k_1 \neq k_2$, (b) $l_1 \neq l_2$.

In case (a), without lost of generality, let $k_1 > k_2$. Then the matrix appears in the first summand of $f^{l_1k_1}_{a_1,..,a_{s_1+k_1-1}}$ has row $y_{k_2}$ without row $y_{k_1}$, and the matrix that appears in the first summand of $f^{l_2k_2}_{b_1,\ldots,b_{s_1+k_2-1}}$ has row $y_{k_1}$without $y_{k_2}$. Consider $m_{12}$, $m_{21}$, $M_{12}$ and $M_{21}$ as defined in Definition 4.2. Assume $M_{12}$ has columns $c_1, \ldots, c_r$ and $M_{21}$ has columns $d_1, \ldots., d_w$. Define $\overline{f^{l_1,k_1}_{a_1,\ldots,a_{s_1+k_1-1},d_1,\ldots,d_w}}$ and $\overline{f^{l_2,k_2}_{b_1,\ldots,b_{s_1+k_2-1},c_1,\ldots,c_r}}$ as in the Definition 4.5. Let $\{c_1, \ldots, c_r, b_1, \ldots, b_{s_1+k_2-1}\} = \{d_1, \ldots, d_w, a_1, \ldots, a_{s_1+k_1-1}\} = \mathcal{I}$, from the way we define $M_{12}$ and $M_{21}$ we have the initial term of $\overline{f^{l_1,k_1}_{\mathcal{I}}}$ is $\text{in}(M_{12}f^{l_1,k_1}_{a_1,\ldots,a_{s_1+k_1-1}})$ and similarly for $\overline{f^{l_2,k_2}_{\mathcal{I}}}$. We will like to apply Lemma 4.6 to this case.

Rewrite $\overline{f^{l_2,k_2}_{\mathcal{I}}}$ as $\alpha_1$:

$$\alpha_1 := \sum_{r=k_2}^{s_2} (-1)^{r+1} \left| \begin{bmatrix} \overline{M_{12}}' \\ Y^{k_2,k_2} \\ Z^{l_2,k_2-1} \\ Z^{r,r} \\ X^{1,l_2-1} \\ Y^{1,r-1} \\ Y^{r+1} \end{bmatrix}_{\mathcal{I}} \right| + \sum_{r=k_2}^{s_2} (-1)^{r+1} \sum_{u=r+1}^{s_1} \left| \begin{bmatrix} \overline{M_{12}}' \\ Y^{k_2,k_2} \\ Z^{l_2,k_2-1} \\ X^{r,r} - Y^{r,r} \\ X^{1,l_2-1} \\ Y^{1,r-1} \\ Y^{r+1,u-1} \\ Z^{u,u} \\ X^{u+1,s_1} \end{bmatrix}_{\mathcal{I}} \right|.$$

Notice that in the first sum of $\alpha_1$, when $r > k_2$, the matrices have repeated row $y_{k_2}$. Hence the determinants are zero. The first sum becomes $\alpha_{11}$:

$$\alpha_{11} := \left| \begin{bmatrix} \overline{M_{12}}' \\ Y^{k_2,k_2} \\ Z^{l_2,k_2-1} \\ Z^{k_2,k_2} \\ X^{1,l_2-1} \\ Y^{1,k_2-1} \\ Y^{k_2+1} \end{bmatrix}_{\mathcal{I}} \right|$$

We notice the leading term of $\alpha_{11}$ is $m_{12}(\text{in} f^{l_2k_2}_{b_1,\dots,b_{s_1+k-1}})$. Let the second sum of $\alpha_1$ be $\alpha_{12}$:

$$\alpha_{12} := \sum_{r=k_2}^{s_2} (-1)^{r+1} \sum_{u=r+1}^{s_1} \left| \begin{bmatrix} \overline{M_{12}}' \\ Y^{k_2,k_2} \\ Z^{l_2,k_2-1} \\ X^{r,r} - Y^{r,r} \\ X^{1,l_2-1} \\ Y^{1,r-1} \\ Y^{r+1,u-1} \\ Z^{u,u} \\ X^{u+1,s_1} \end{bmatrix}_{\mathcal{I}} \right|.$$

Lemma 4.6 provided if $\alpha_{12}$ is a combination of elements of $\mathcal{G}$ such that the leading term of each summand is smaller than $m_{12} f^{l_2k_2}_{b_1,\dots,b_{s_1+k_2-1}}$. Observe that in the sum of $\alpha_{12}$, when $r > k_2$, the matrices have repeated row $y_{k_2}$. Hence their determinants are zero.

We are only left with $r = k_2$, and $\alpha_{12}$ becomes

$$\alpha_{13} := \quad (-1)^{k_2+1} \sum_{u=k_2+1}^{s_1} \left| \begin{bmatrix} \overline{M_{12}}' \\ Y^{k_2,k_2} \\ Z^{l_2,k_2-1} \\ X^{k_2,k_2} - Y^{k_2,k_2} \\ X^{1,l_2-1} \\ Y^{1,k_2-1} \\ Y^{k_2+1,u-1} \\ Z^{u,u} \\ X^{u+1,s_1} \end{bmatrix}_{\mathcal{I}} \right|.$$

We apply Lemma 3.3 on $\alpha_{13}$, then $\alpha_{13}$ becomes $\alpha_{14}$:

$$\alpha_{14} := \quad (-1)^{k_2+1} \sum_{u=k_2+1}^{s_1} \left| \begin{bmatrix} \overline{M_{12}}' \\ Y^{k_2,k_2} \\ X^{l_2,l_2} - Y^{l_2,l_2} \\ Z^{l_2+1,k_2-1} \\ Z^{k_2,k_2} \\ X^{1,l_2-1} \\ Y^{1,k_2-1} \\ Y^{k_2+1,u-1} \\ Z^{u,u} \\ X^{u+1,s_1} \end{bmatrix}_{\mathcal{I}} \right| + (-1)^{k_2+1} \sum_{u=k_2+1}^{s_1}$$

$$\sum_{\{p_1,p_2,q_1,\ldots,q_{s_1+k_2-1}\}=\mathcal{I}} \left( \pm(g_{l_2p_1,k_2,p_2} - g_{l_2p_2,k_2p_1}) \left| \begin{bmatrix} \overline{M_{12}}' \\ Y^{k_2,k_2} \\ Z^{l_2+1,k_2-1} \\ X^{1,l_2-1} \\ Y^{1,k_2-1} \\ Y^{k_2+1,u-1} \\ Z^{u,u} \\ X^{u+1,s_1} \end{bmatrix}_{q_1,\ldots,q_{s_1+k-2}} \right| \right).$$

After removing the repeated row $y_{l_2}$ in the first sum in the above expression for $\alpha_{14}$, let this sum be $\alpha_{15}$:

$$\alpha_{15}: \quad = (-1)^{k_2+1}\sum_{u=k_2+1}^{s_1}\left|\begin{bmatrix} \overline{M_{12}}' \\ Y^{k_2,k_2} \\ X^{l_2,l_2} \\ Z^{l_2+1,k_2-1} \\ Z^{k_2,k_2} \\ X^{1,l_2-1} \\ Y^{1,k_2-1} \\ Y^{k_2+1,u-1} \\ Z^{u,u} \\ X^{u+1,s_1} \end{bmatrix}_{\mathcal{I}}\right| = \pm\sum_{u=k_2+1}^{s_1}(-1)^{u+1}\left|\begin{bmatrix} \overline{M_{12}}' \\ Z^{l_2+1,k_2} \\ Z^{u,u} \\ X^{1,l_2} \\ Y^{1,u-1} \\ X^{u+1,s_1} \end{bmatrix}_{\mathcal{I}}\right|$$

Now $\alpha_{15}$ becomes

$$\begin{aligned}\alpha_{16} := & \sum_{\{p_1,\dots,p_{v-1},q_1,\dots,q_{s_{1+k_2}}\}=\mathcal{I}} \pm|M^{12}_{p_1,\dots,p_{v-1}}| \left(\sum_{u=k_2+1}^{s_1}(-1)^{u+1}\left|\begin{bmatrix} Z^{l_2+1,k_2} \\ Z^{u,u} \\ X^{1,l_2} \\ Y^{1,u-1} \\ X^{u+1,s_1} \end{bmatrix}_{\mathcal{I}}\right|\right) \\ = & \sum_{\{p_1,\dots,p_{v-1},q_1,\dots,q_{s_{1+k_2}}\}=\mathcal{I}} \pm|M^{12}_{p_1,\dots,p_{v-1}}|p^{l_2+1,k_2+1}_{q_1,\dots,q_{s_1+k_2}}.\end{aligned}$$

Here $\{p^{l_2+1,k_2+1}_{q_1,\dots,q_{s_1+k_1}}\}$ are as defined in lemma 4.7, and the proof of lemma 4.7 shows that they are in $\mathcal{L}$ . This shows $\alpha_{12}$ is a combination of elements of $\mathcal{G}$ such that the leading term of each summand is smaller than $m_{12}f^{l_2k_2}_{b_1,\dots,b_{s_1+k_2-1}}$.

We can do the same to $\overline{f^{l_1,k_1}_{\mathcal{I}}}$ and show the second part of the sum of $\overline{f^{l_1,k_1}_{\mathcal{I}}}$ is a combination of elements of $\mathcal{G}$ such that the leading term of each summand is smaller than $\mathrm{in}(m_{21}f^{l_1k_1}_{a_1,\dots,a_{s_1+k_1-1}})$.

In case (b): assume $k_1 = k_2$ and $l_1 < l_2 \le k_2 = k_1$. The proof technique is very similar to case (a). Notice that the first matrix appearing in the expression for $f^{l_1k_1}_{a_1,\dots,a_{s_1+k_1-1}}$ has row $z_{l_1}$ without row $x_{l_1}$ and the first matrix appearing in the expression for $f^{l_2k_2}_{b_1,\dots,b_{s_1+k_1-1}}$ has row $x_{l_1}$ without row $z_{l_1}$. Since $l_1 \le l_2 - 1$ and $l_1 \le k_2 - 1$, each matrix of $\overline{f^{l_2,k_2}_{b_1,\dots,b_{s_1+k_2-1},c_1,\dots,c_r}}$ has the rows $x_{l_1}$ and $y_{l1}$. They also all have row $z_{l_1}$. Applying lemma 3.2 gives all the determinants of those matrices are in $(\{g_{l_1i,l_1j}\})$. □

**Lemma 4.31.** *$h_{PQ} = 0$ if $P$, $Q \in \{f^{l,k}_{a_1,\dots,a_{s_1+k-1}}, U_{p_1,q_1,a_1,\dots,a_{s_1}}$ ,
$W_{p,q,a_1,\dots,a_p,a_{s_2+1},\dots,a_{s_1},b_1,\dots,b_{s_2}}$,
$W^{p_1+1,v}_{p_1,q_1,a_1,\dots,a_{p1},a_{s_2+1},\dots,a_{s_1},b_1,\dots,b_{p_1+1},b_{p_1+2},b_{p_1+3}.,b_{s_2},b'_{p_1+1},b'_{p_1+2},\dots,b'_v}$,*

$V_{p_l,\dots,p_k,a_1,\dots,a_{k-1},b_1,\dots,b_{s_1}}$, $V^{k,w}_{p_l,\dots,p_k,a_1,\dots,a_{k-1},b_1,\dots,b_{s_1},b'_k,b'_{k+1},\dots,b'_w}$, $H^{l,k,q}_{a_1,\dots,a_{s_1+k-1}}$, $I^{l,k,q}_{a_{s_1+k-1},\dots,a_{l+s_2-1},a_{l-1},\dots,a_1,b_1,\dots,b_{s_2}}$, ${}^{k,w}I^{l,k,q}_{q_l,\dots,q_k,b_{s_1},\dots,b_{s_2},\dots,b_1,a_{l-1},\dots,a_1,b'_k,\dots,b'_w}\}$.

*Proof.* For the purpose of the proof, we drop the column indices. The remainders of $S$-pair of $f^{l,k}$ and $U$ are in the ideal generated by $(\{V\},\{W\},\{g\},\{Y\})$. Similarly, the remainders of $S$-pair of $f^{l,k}$ and $W$ are in $(\{V^{k,w}\}, \{W^{p_1+1,v}\},\{g\},\{Y\})$, and the remainders of $S$-pair of $f^{l,k}$ and $W^{p_1+1,v}$ are in $(\{V^{k,v+1}\}, \{W^{p_1+1,v+1}\}, \{g\}, \{Y\})$. The remainders of $S$-pair of $f^{l,k}$ and $V$ are $V^{k,w}$ and the remainders of $S$-pair of $f^{l,k}$ and $V^{k,w}$ are $V^{k,w+1}$. The remainders of $S$-pair of $f^{l,k}$ and $H^{l,k,q}$ are in $(\{I^{l,k,q}\}, \{g\}, \{Y\})$ and the remainders of $S$-pair of $f^{l,k}$ and $I^{l,k,q}$ are in $(\{{}^{k,w}I^{l,k,q}\}, \{g\}, \{Y\})$. Finally the remainders of $S$-pair of $f^{l,k}$ and ${}^{k,w}I^{l,k,q}$ are in $(\{{}^{k,w+1}I^{l,k,q}\}, \{g\}, \{Y\})$. All the other $S$-pair of elements have similar relationship as above. □

We complete the proof of Theorem 4.25.

*Proof of Lemma 2.7:* Notice that $x_{11}$ is a non zero-divisor on $k[X,Y,Z]/\mathrm{in}(\mathcal{L})$. Since the only possible elements of $\mathcal{G}$ that leading monomials are divisible by $x_{11}$ are $U_{1,1,1,a_2,\dots,a_{s_1}}$, $W_{1,1,1,a_{s_2+1},\dots,a_{s_1},b_1,\dots,b_{s_2}}$ and $W^{2,v}_{1,1,1,a_{s_2+1},\dots,a_{s_1},b_1,\dots,b_{s_2},b'_2,\dots,b'_v}$. But those monomials are divisible by the leading monomials of $f^{1,1}_{1,a_2,\dots,a_{s_1}}$, $V_{1,b_1,\dots,b_{s_2},a_{s_2+1},\dots,a_{s_1}}$ and $V^{2,v}_{1,b_1,\dots,b_{s_2},a_{s_2+1},\dots,a_{s_1},b'_2,\dots,b'_v}$. Hence $x_{11}$ is also a non zero-divisor on $k[X,Y,Z]/\mathcal{L}$. □

The following example is computed in Singular [GPS]. This example gives us an idea what does the initial ideal looks like.

**Example 4.32.** Let $X$, $Y$, $Z$ be a $3\times 4$ matrices, $X_{3,4}$, $Y_{2,4}$ are $3\times 4$ and $2\times 4$ submatrices of $X$ and $Y$ then the defining ideal of the $\mathcal{R}(\mathbb{D})$ is generated by $I_3(X_{3,4})$, $I_2(Y_{2,4})$, $g_{ij,lk}$ where $1\le i,l\le 3$, $1\le l,k\le 4$ and

$$f_{a_1,\dots,a_3} = \begin{vmatrix} z_{1a_1} & z_{1a_2} & z_{1a_3} \\ x_{2a_1} & x_{2a_2} & x_{2a_3} \\ x_{3a_1} & x_{3a_2} & x_{3a_3} \end{vmatrix} + \begin{vmatrix} y_{1a_1} & y_{1a_2} & y_{1a_3} \\ z_{2a_1} & z_{2a_2} & z_{2a_3} \\ x_{3a_1} & x_{3a_2} & x_{3a_3} \end{vmatrix},$$

where $1\le a_1<a_2<a_3\le 4$. The initial ideal of $\mathcal{L}$ via the term order defined in Theorem 4.25 is generated by $\{x_{1a_3}x_{2a_2}x_{3a_1}\}_{1\le a_1<a_2<a_3\le 4}$, $\{y_{1b_2}y_{2b_1}\}_{1\le b_1<b_2\le 4}$, $\{z_{ij}x_{lk}\}_{i<l}$ or $_{i=l}$ and $j<k$, $\{z_{1a_1}y_{2a_2}y_{3a_3}\}_{1\le a_1<a_3<a_2\le 4}$, $z_{11}z_{22}y_{34}y_{33}$, $z_{21}x_{14}y_{13}y_{32}$, $z_{12}z_{21}x_{12}y_{14}y_{33}$, $z_{13}z_{21}x_{13}y_{14}y_{32}$, $z_{31}x_{14}x_{23}y_{32}$, $\{z_{2j}x_{1a_3}x_{2a_2}y_{3a_1}\}_{1\le a_1<a_2\le j<a_3\le 4}$, or $_{1\le a_2\le j<a_1<a_3\le 4}$, $\{z_{2j}x_{1a_3}y_{2a_2}y_{1a_1}\}_{1\le a_1<a_2<a_3,a_2<j}$, $\{z_{1j}x_{1a_3}y_{2a_2}y_{3a_1}\}_{1\le a_1<a_2<a_3\le j\le 4}$, or $_{1\le a_1<a_3\le j<a_2\le 4}$, $\{z_{1j}x_{1a_3}y_{1b_1}y_{2b_2}y_{3b_3}\}_{1\le b_2<b_3<a_3\le j\le 4}$, or $_{1\le a_1<a_3\le j<a_2\le 4}$. We can see the variable $x_{11}$ is not in the generating set of the initial ideal of $\mathcal{L}$.

## References

[BH] W. Bruns and J. Herzog, Cohen-Macaulay rings, Cambridge Studies in Advanced Mathematics 39, Cambridge University Press, Cambridge, 1993.

[E] Commutative algebra with a view toward algebraic geometry, Graduate Texts in Mathematics 150, Springer-Verlag, New York, 1995.

[GS] D. Grayson and M. Stillman, Macaulay 2, A computer algebra system for computing in algebraic geometry and commutative algebra, available through anonymous ftp from http://www.math.uiuc.edu/Macaulay2.

[GPS] G. Greuel, G. Pfister, and H. Schonemann, Singular 3.1.0, A computer algebra system for polynomial computations, available through http://www.singular.uni-kl.de.

[HSV1] J. Herzog, A. Simis and W. Vasconcelos,Approximation complexes of blowing-up rings. J. Algebra 74 (1982), 466–493.

[HSV2] J. Herzog, A. Simis and W. Vasconcelos, Approximation complexes of blowing-up rings. II. J. Algebra 82 (1983), 53–83.

[HTZ] J. Herzog, Z. Tang and S. Zarzuela, Symmetric and Rees algebras of Koszul cycles and their Groebner bases, Manuscripta Math. 112 (2003), 489-509.

[HVV] J. Herzog, W. Vasconcelos and R. Villarreal, Ideals with sliding depth. Nagoya Math. J. 99 (1985), 159–172.

[H1] C. Huneke, On the symmetric and Rees algebra of an ideal generated by a $d$-sequence. J. Algebra 62, 268-275 (1980).

[H2] C. Huneke, Determinantal ideals of linear type. Arch. Math. (Basel) 47 (1986), 324–329.

[M] A. Micali, Sur les algebres universelles. Ann. Institut Fourier, 14, 33-88 (1964).

[SSU] A. Simis, K. Smith and B. Ulrich, An algebraic proof of Zak's inequality for the dimension of the Gauss image, Math. Z. 241 (2002), 871-881.

[SU] A. Simis and B. Ulrich, On the ideal of an embedded join, J. Algebra 226 (2000), 1-14.

[SS] B. Sturmfels and S. Sullivant, Combinatorial secant varieties, Pure Appl. Math. 2 (2006), 867-891.

[V] G. Valla, On the symmetric and Rees algebras of an ideal. Manuscripta Math. 30, 239-255 (1980).

[Vi] R. H. Villarreal, Cohen-Macaulay graphs. Manuscripta Math. 66 (1990), 277–293.

Department of Mathematics,

University of California, Riverside, CA 92521, USA

e-mail: linkuei@ucr.edu